




openheart Association of HFA-PEFF score with clinical outcomes after catheter ablation for atrial fibrillation

Taiji Okada ¹, Takeshi Kitai,² Atsushi Kobori,³ Madoka Sano,¹ Ryosuke Murai,¹ Toshiaki Toyota ¹, Yasuhiro Sasaki,¹ Tomohiko Taniguchi ^{1,4,5}, Kitae Kim,¹ Natsuhiko Ehara,³ Makoto Kinoshita,¹ Yutaka Furukawa¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/openhrt-2023-002526>).

To cite: Okada T, Kitai T, Kobori A, *et al.* Association of HFA-PEFF score with clinical outcomes after catheter ablation for atrial fibrillation. *Open Heart* 2024;**11**:e002526. doi:10.1136/openhrt-2023-002526

Received 14 October 2023
Accepted 3 January 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Cardiovascular Medicine, Kobe City Medical Center General Hospital, Kobe, Japan

²Department of Cardiovascular Medicine, National Cerebral and Cardiovascular Center, Suita, Japan

³Cardiology, Kobe City Medical Center General Hospital, Kobe, Japan

⁴Kobe City Medical Center General Hospital, Kobe, Japan

⁵Cardiology, Kokura Memorial Hospital, Kitakyushu, Japan

Correspondence to

Dr Taiji Okada; taiji_okada@kcho.jp

ABSTRACT

Background The Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology (HFA-PEFF) score has been developed for diagnosing heart failure with preserved ejection fraction (HFpEF), which is frequently associated with atrial fibrillation (AF). We aimed to investigate whether preprocedural HFA-PEFF score could be used to predict clinical outcomes in patients with AF who underwent catheter ablation (CA).

Methods Overall, 1679 patients with AF who underwent primary CA (71±10 years, 1218 males (72.5%), median follow-up duration 3.3 years) from July 2011 to December 2019 were included in this retrospective study. HFpEF was defined as an HFA-PEFF score ≥5. The primary study outcome was 5-year major adverse cardiovascular and cerebrovascular events (MACCE), which is a composite of all-cause death, hospitalisation for heart failure (HF) and hospitalisation for stroke.

Results The prevalence of HFpEF was 32.3%, but only 7.7% were diagnosed with HF at the time of CHADS₂ scoring. Five-year MACCE occurred in 77 patients (4.6%). The cumulative 5-year incidence of MACCE was significantly higher in the HFpEF group than in the non-HFpEF group (11.2% vs 4.8% at 5 years, p<0.001). In the multivariable analysis, HFpEF by the HFA-PEFF score was associated with MACCE (adjusted HR 1.65, 95% CI 1.02 to 2.65, p=0.041).

Conclusions Early detection of HFpEF using the HFA-PEFF score may have clinical applications in guiding therapeutic decision-making and improving prognosis by preventing HF and stroke in patients with AF undergoing CA.

INTRODUCTION

Atrial fibrillation (AF) and heart failure with preserved ejection fraction (HFpEF) are common conditions with increasing prevalence and are associated with increased morbidity and mortality.^{1 2} However, the diagnosis of HFpEF remains challenging.³ HFpEF is defined as symptomatic heart failure (HF) with a left ventricular ejection fraction (LVEF) ≥50% according to a universal definition,⁴ and two other scores have been

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Atrial fibrillation (AF) and heart failure (HF) with preserved ejection fraction (HFpEF) are common conditions with increasing prevalence and are associated with increased morbidity and mortality.
- ⇒ The diagnosis of HFpEF remains challenging, and differences in clinical events after catheter ablation (CA) between HFpEF and non-HFpEF have been variously reported.

WHAT THIS STUDY ADDS

- ⇒ In patients with AF undergoing CA, using the Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology (HFA-PEFF) score could help identify more patients with HFpEF, which were initially identified as having HF.
- ⇒ In addition, HFpEF diagnosed by the HFA-PEFF score is associated with a higher risk of hospitalisation for HF, hospitalisation for stroke and death (major adverse cardiovascular and cerebrovascular events) after CA.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ In patients undergoing CA for AF, using the HFA-PEFF score may help to identify patients with underdiagnosed HFpEF, which could, in turn, improve their prognosis.
- ⇒ Future studies using the HFA-PEFF and the H₂FPEF scores could detect 'early HFpEF' and may represent a potential target for CA, which may contribute to improved prognosis through the prevention of HF and stroke.

proposed.^{5 6} The Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology (HFA-PEFF) score is one of the steps in the algorithm developed by the Heart Failure Association of the European Society of Cardiology, based on echocardiographic and laboratory findings,⁶ and the H₂FPEF score includes six demographic, clinical and echocardiographic

variables.⁵ These scores have shown different diagnostic performance^{7,8} and outcome prediction^{9,10} in different cohorts.

Catheter ablation (CA) is a therapeutic strategy for patients with AF in the rhythm control strategy. CA is a reasonable treatment for patients with AF, particularly when the symptoms are not adequately controlled or tolerated with pharmacological therapy. Some studies have shown that the restoration of sinus rhythm by CA for treating HFpEF can reduce clinical events.^{11,12} However, a large retrospective cohort study¹³ and the Catheter Ablation versus Antiarrhythmic Drug Therapy for Atrial Fibrillation trial,¹⁴ in which most participants had preserved LVEF, did not demonstrate a beneficial effect on the primary outcome. One reason for this may be that HFpEF was variously defined as LVEF alone. Therefore, we hypothesised that using the HFA-PEFF score could help identify patients with AF at high risk of death and hospitalisation for HF and stroke.

In this study, we performed a retrospective analysis of patients with AF undergoing CA to assess whether the diagnosis of HFpEF using preprocedural HFA-PEFF score is associated with clinical outcomes.

METHODS

Study population

From July 2011 to December 2019, 2123 consecutive patients with AF who underwent primary CA were screened. We excluded patients with prior cardiac

surgery, prior catheter interventions for valve disease, haemodialysis, significant left-side valvular disease (defined as moderate or severe mitral stenosis, moderate or severe mitral regurgitation, moderate or severe aortic regurgitation and moderate or severe aortic stenosis),¹⁵ atrial septal defects, LVEF <50% and hypertrophic cardiomyopathy. Finally, 1679 patients were analysed retrospectively in this study (figure 1).

Echocardiographic measurements

Two-dimensional echocardiography was performed before CA using the Aplio Artida (Canon Medical Systems, Otawara, Japan) or Epic 7, iE 33 (Philips Healthcare, Eindhoven, The Netherlands) ultrasound systems, and data were obtained following the recommendations of the American Society of Echocardiography and the European Association of Cardiovascular Imaging.^{16,17} All measurements were quantified as an average of three consecutive heartbeats for patients with sinus rhythm and as an index beat for patients with AF.¹⁸

HFA-PEFF and H₂FPEF scores

The HFA-PEFF score was calculated as the sum of (i) the functional domain (e' , E/e' and tricuspid regurgitation peak velocity (TRV) excluding global longitudinal strain), (ii) the morphological domain (left atrial volume index, left ventricular mass index, relative wall thickness or LV end-diastolic wall thickness) and (iii) the laboratory domain (N-terminal




Figure 1 Flow chart illustrating the study population. Overall, 2123 patients with AF who underwent primary ablation were screened, and finally, 1679 patients were analysed retrospectively in this study. The circles in the diagram represent the percentage of heart failure with preserved left ventricular ejection fraction at each score. AF, atrial fibrillation; HFA-PEFF, Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology; HFpEF, heart failure preserved left ventricular ejection fraction; LVEF, left ventricular ejection fraction.

Table 1 Characteristics and outcome of the study population divided by HFpEF or non-HFpEF in the HFA-PEFF score

	All (n=1679)	HFpEF (n=542)	Non-HFpEF (n=1137)	P value*
Demographic data				
Age, years	71±10	76±7	68±11	<0.001
Sex, male	1218 (72.5)	325 (60.0)	893 (78.5)	<0.001
Height, m	1.66±0.09	1.62±0.09	1.67±0.09	<0.001
Weight, kg	65.6±12.4	62.5±12.4	67.1±12.1	<0.001
Body mass index, kg/m ²	23.8±3.5	23.7±3.6	23.9±3.5	0.128
Body surface area, m ²	1.72±0.19	1.66±0.19	1.75±0.18	<0.001
CHADS ₂ score	1 (1–2)	2 (1–2)	1 (0–2)	<0.001
Congestive heart failure	130 (7.7)	72 (13.3)	58 (5.1)	<0.001
Hypertension	925 (55.1)	345 (63.7)	580 (51.0)	<0.001
Age ≥75 years	664 (39.5)	310 (57.2)	354 (31.1)	<0.001
Diabetes mellitus	245 (14.6)	85 (15.7)	160 (14.1)	0.382
Prior stroke	147 (8.8)	66 (12.2)	81 (7.1)	0.001
Coronary artery disease	142 (8.5)	53 (9.9)	89 (7.9)	0.178
Chronic kidney disease ≥grade 3	587 (45.7)	247 (45.7)	340 (30.1)	<0.001
Paroxysmal AF	1085 (64.6)	343 (63.3)	742 (65.3)	0.429
2D echocardiographic data				
LV end-diastolic dimension, mm	45.6±4.5	45.6±4.8	45.7±4.4	0.703
LVEF, %	62.4±4.0	62.5±4.2	62.3±3.9	0.301
LV mass index, g/m ²	82.4±17.0	87.8±18.3	79.8±15.7	<0.001
LA diameter, mm	38.2±6.4	40.4±6.0	37.2±6.3	<0.001
LA volume index, mL/m ²	43.2±14.4	50.2±13.2	40.0±14.6	<0.001
LA emptying fraction, %	32.6±14.7	29.6±14.6	34.0±14.6	<0.001
E velocity, cm/s	75.4±20.8	80.9±21.4	72.8±18.4	<0.001
Septal e', cm/s	7.9±2.3	6.8±1.7	8.4±2.4	<0.001
Lateral e', cm/s	10.1±3.0	8.7±2.2	10.7±3.0	<0.001
Mean E/e'	9.0±3.0	10.9±3.2	8.0±2.4	<0.001
TRV, m/s (n=1547)	2.3±0.3	2.5±0.3	2.3±0.3	<0.001
Moderate or severe TR	161 (9.6)	83 (15.3)	78 (6.9)	<0.001
Laboratory data				
BNP, pg/mL (n=826)	50.0 (23.4–101.5)	88.8 (53.0–154.7)	32.6 (16.7–73.0)	<0.001
NT pro-BNP, pg/mL (n=942)	192.7 (71.3–622.8)	602.5 (240.8–1035.8)	100.5 (44.7–347.8)	<0.001
Estimated GFR, mL/min/1.73 m ²	2.4±0.3	61.9±14.7	67.4±14.3	<0.001
HFA-PEFF score	4 (3–5)	5 (5–6)	3 (2–4)	<0.001
Outcome				
5-year MACCE	77 (4.6)	42 (7.7)	35 (3.1)	<0.001
5-year death	38 (2.3)	20 (3.7)	18 (1.6)	0.007
5-year hospitalisation for heart failure	11 (0.7)	9 (1.7)	2 (0.2)	<0.001
5-year hospitalisation for stroke	34 (2.0)	18 (3.3)	16 (1.4)	0.009
5-year AF recurrence	495 (29.5)	189 (34.9)	306 (26.9)	0.001
Data are expressed as mean±SD or median (IQR) or number (%).				
*P values refer to the difference between HFpEF and non-HFpEF.				
AF, atrial fibrillation; BNP, B-type natriuretic peptide; 2D, two-dimensional; GFR, glomerular filtration rate; HFA-PEFF, Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology; HFpEF, heart failure preserved left ventricular ejection fraction; LA, left atrial; LV, left ventricular; LVEF, left ventricular ejection fraction; MACCE, major adverse cardiovascular and cerebrovascular events; NT pro-BNP, N-terminal prohormone brain natriuretic peptide; TR, tricuspid regurgitation; TRV, tricuspid regurgitation peak velocity.				

Table 2 Difference in factors of the HFA-PEFF score

	All (n=1679)	HFpEF (n=542)	Non-HFpEF (n=1137)	P value*
HFA-PEFF score	4 (3–5)	5 (5–6)	3 (2–4)	<0.001
Functional				
Mean E/e' 9–14, 1 pt.	650 (39.2)	330 (60.9)	320 (28.7)	<0.001
Mean E/e' ≥15, 2 pts.	89 (5.4)	70 (12.9)	19 (1.7)	<0.001
Septal e' <7 or Lateral e' <10, 2 pts.	960 (57.8)	475 (87.6)	485 (43.4)	<0.001
TRV ≥2.8 m/s (n=1547), 2 pts.	89 (5.8)	64 (12.4)	25 (2.4)	<0.001
Morphological				
LAVI 29–34 mL/m ² , 1 pt.	233 (13.3)	12 (2.2)	211 (18.6)	<0.001
LAVI >34 mL/m ² , 2 pts.	1203 (71.7)	519 (95.8)	684 (60.2)	<0.001
LVMI >115/95 g/m ² (M/F), 1 pt.	127 (7.6)	80 (14.8)	47 (4.1)	<0.001
RWT >0.42, 1 pt.	624 (37.2)	237 (43.7)	387 (34.0)	<0.001
LVMI ≥149/122 g/m ² (M/F) and RWT >0.42, 2 pts.	3 (0.2)	3 (0.2)	0 (0.0)	0.012
LV wall thickness ≥12 mm, 1 pt.	68 (4.1)	42 (7.7)	26 (2.3)	<0.001
Biomarker				
BNP 35–80 or NT-pro BNP 125–220 at sinus rhythm, BNP 105–240 or NT-pro BNP 365–660 at AF, 1 pt.	466 (27.8)	217 (40.0)	249 (21.9)	<0.001
BNP >80 or NT-pro BNP >220 at sinus rhythm, BNP >240 or NT-pro BNP >660 at AF, 2 pts.	446 (26.6)	325 (60.0)	121 (10.6)	<0.001

Data are expressed as mean±SD or median (IQR) or number (%).

*P values refer to the difference between HFpEF and non-HFpEF.

AF, atrial fibrillation; BMI, body mass index; BNP, B-type natriuretic peptide; F, female; HFA-PEFF, Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology; HFpEF, heart failure preserved left ventricular ejection fraction; LA, left atrial; LAVI, left atrial volume index; LV, left ventricular; LVEF, left ventricular ejection fraction; LVMI, left ventricular mass index; M, male; NT pro-BNP, N-terminal prohormone brain natriuretic peptide; pt., patient; RWT, relative wall thickness; TRV, tricuspid regurgitation peak velocity.

prohormone brain natriuretic peptide (BNP) or BNP). A major criterion scores 2 points and a minor criterion scores 1 point within each domain (online supplemental figure S1A).

The H₂FPEF score is used with some modifications and calculated as the sum of (i) AF, 3 points; (ii) obesity defined as body mass index >30 kg/m², 2 points; (iii) age >60 years, 1 point; (iv) hypertension diagnosis or treatment with two or more antihypertensive drugs, 1 point; (v) E/e' ratio >9, 1 point and (vi) TRV >2.8 m/s, 1 point (online supplemental figure S2A).

In this study, HFpEF was defined as an HFA-PEFF score ≥5.

Study outcomes

The primary study outcome was 5-year major adverse cardiovascular and cerebrovascular events (MACCE), which was a composite of all-cause death, hospitalisation for HF and hospitalisation for stroke. The secondary study outcomes were 5-year all-cause death, 5-year hospitalisation for HF and 5-year hospitalisation for stroke. All adverse events were collected during follow-up using electronic records, laboratory findings and echocardiographic reports.

Statistical analysis

Categorical data are expressed as numbers and percentages, while quantitative data are described as the mean±SD or median and IQR. The comparison between patients with HFpEF and those without HFpEF was performed using the unpaired t-test or Mann-Whitney U test, according to the normality of the data. The χ^2 test, or Fisher's exact test, was performed to compare categorical data. For four-group comparisons of baseline characteristics, we used analysis of variance with Dunnett's test as a post hoc analysis. Survival curves were obtained using the Kaplan-Meier method and compared using the log-rank test. The unadjusted and adjusted HRs and 95% CIs for the associations of the clinical and echocardiographic variables and HFpEF with the primary end point were obtained from univariate and multivariate Cox regression analysis. In the multivariate Cox regression, model 1 included HFpEF, age and sex, and model 2 included model 1, hypertension, diabetes mellitus, prior stroke, chronic kidney disease ≥grade 3 and moderate or severe tricuspid regurgitation (TR).




Figure 2 Kaplan-Meier analyses show the cumulative rate of the primary composite end point (A), all-cause death (B), hospitalisation for HF (C) and hospitalisation for stroke (D) stratified by the Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology score. HFpEF, heart failure preserved left ventricular ejection fraction.

Subgroup analysis for MACCE was repeated in patients with paroxysmal AF, persistent AF and AF recurrence, defined as any documented atrial arrhythmia of >30 s after CA, and without AF recurrence using the Kaplan-Meier method and compared using the log-rank test. The sensitivity analysis for MACCE was performed using a multivariate Cox regression analysis including all predictors of MACCE with p values <0.10 in the univariate analysis, with

stepwise backward regression using a probability to leave of 0.10. In addition, multivariate Cox regression analysis performed by replacing HFpEF by HFAPEFF score with HFpEF by H_2 FPEF score, which was defined as an H_2 FPEF score ≥ 6 .

P values <0.05 were considered statistically significant. The statistical analysis was performed using SPSS Statistics V.23.0 (IBM, Armonk, New York, USA).

Table 3 Univariable and multivariable analyses for MACCE for the HFA-PEFF score

	Univariate analysis		Multivariate analysis 1		Multivariate analysis 2	
	HR (95% CI)	P value	HR (95% CI)	P value	HR (95% CI)	P value
HFpEF by the HFA-PEFF score	2.56 (1.63 to 4.01)	<0.001	1.75 (1.09 to 2.80)	0.021	1.65 (1.02 to 2.65)	0.041
Age	1.09 (1.05 to 1.12)	<0.001	1.08 (1.04 to 1.11)	0.843	1.07 (1.03 to 1.10)	<0.001
Male	0.73 (0.46 to 1.18)	0.199	1.08 (0.66 to 1.76)	0.762	1.13 (0.69 to 1.85)	0.630
Hypertension	1.39 (0.87 to 2.21)	0.165			1.01 (0.63 to 1.62)	0.970
Diabetes mellitus	0.97 (0.51 to 1.83)	0.922			0.95 (0.50 to 1.79)	0.862
Prior stroke	1.57 (0.81 to 3.04)	0.186			1.30 (0.66 to 2.55)	0.444
Chronic kidney disease \geq grade 3	1.95 (1.25 to 3.04)	0.003			1.35 (0.85 to 2.14)	0.202
Moderate or severe TR	2.80 (1.65 to 4.75)	<0.001			1.87 (1.09 to 3.22)	0.024

Multivariate analysis model 1 includes HFpEF, age and sex. Multivariate analysis model 2 includes model 1, hypertension, diabetes mellitus, prior stroke, chronic kidney disease defined as estimated GFR <60 mL/min/1.73 m² and moderate or severe TR. GFR, glomerular filtration rate; HFA-PEFF, Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology; HFpEF, heart failure with preserved left ventricular ejection fraction; MACCE, major adverse cardiovascular and cerebrovascular events; TR, tricuspid regurgitation.

RESULTS

Patient characteristics

The study population included 1679 patients, with a mean age of 71 ± 10 years, including 1218 males (72.5%); the median HFA-PEFF score was 4 (IQR 3–5, online supplemental figure S1B) and HFpEF was diagnosed in 542 patients (32.3%). The baseline characteristics are presented in table 1. HFpEF had older patients, more female, lower body surface area, more hypertension, prior stroke, chronic kidney disease, diastolic dysfunction, a higher estimated LV filling pressure by echocardiographic findings and no differences in AF type compared with the non-HFpEF group. The difference in the factors of the HFA-PEFF score is shown in table 2.

Prognostic value of the HFA-PEFF score

Five-year MACCE occurred in 77 patients (4.6%) (median follow-up duration: 3.3 years (IQR 3.0–5.0)). All-cause death occurred in 38 patients (2.3%), hospitalisation for HF occurred in 11 patients (0.7%) and hospitalisation for stroke occurred in 34 patients (2.0%) within 5 years after CA (table 1). The cumulative 5-year incidence of MACCE was significantly higher in the HFpEF group than in the non-HFpEF group (11.2% vs 4.8% at 5 years, $p < 0.001$, figure 2A) and the cumulative 5-year incidence of all-cause death, hospitalisation for HF and hospitalisation for stroke were also significantly higher in the HFpEF group than in the non-HFpEF group (figure 2B–D).

The results of the univariable and multivariable Cox regression analyses are presented in table 3. HFpEF was associated with MACCE in the univariate analysis (HR 2.56, 95% CI 1.63 to 4.01, $p < 0.001$). Even after adjusting for model 1, which included age and sex, and model 2, which included age, sex, hypertension, diabetes mellitus, prior stroke, chronic kidney disease and moderate or severe TR, HFpEF was associated with MACCE (HR 1.75, 95% CI 1.09 to 2.80, $p = 0.021$, and HR 1.65, 95% CI 1.02 to 2.65, $p = 0.041$, respectively).


Subgroup and sensitivity analysis for MACCE

The results of the subgroup analysis were similar to those of the primary analysis, regardless of prior AF type or 5-year AF recurrences (figure 3). In the univariate analysis, HFpEF, age, body mass index, body surface area, chronic kidney disease \geq grade 3, LV end-diastolic dimension and moderate or severe TR were $p < 0.10$. In the multivariable Cox regression analysis that included these factors, HFpEF, age and moderate or severe TR were independent predictors for MACCE (HR 1.65, 95% CI 1.03 to 2.64; HR 1.07, 95% CI 1.04 to 1.10 and HR 1.83, 95% CI 1.07 to 3.16, respectively) (online supplemental table S1).


Prognostic value of the H₂FPEF score

HFpEF by the H₂FPEF score was observed in 560 patients (33.3%, online supplemental figure S2B). These patients exhibited similar characteristics to those diagnosed with HFpEF by HFA-PEFF score: older patients, more female, lower body surface area, more hypertension, prior stroke, chronic kidney disease, diastolic dysfunction, a higher estimated LV filling pressure by echocardiographic findings and no significant differences in AF type compared with the non-HFpEF group. The features only in HFpEF by the H₂FPEF score were additional features: higher prevalence of diabetes mellitus, coronary artery disease, larger LV dimension and lower LVEF compared with the non-HFpEF group (online supplemental table S2). However, by the H₂FPEF score, the cumulative 5-year incidence of MACCE did not differ significantly between the HFpEF and non-HFpEF groups (8.9% vs 5.9% at 5 years, $p = 0.065$, online supplemental figure S3A), and the cumulative 5-year incidence of hospitalisation for HF and hospitalisation for stroke were also not different between the HFpEF and non-HFpEF groups, although only the cumulative 5-year incidence of all-cause death was significantly higher in the HFpEF group than the non-HFpEF group (5.8% vs


A Paroxysmal AF



B Persistent AF



C 5-year AF recurrence



D 5-year AF no recurrence




Figure 3 Kaplan-Meier analyses show the cumulative rate of the primary composite end point stratified by the HFA-PEFF score in the subgroup of patients with paroxysmal AF (A), persistent AF (B), AF recurrence (C) and AF no recurrence (D). AF, atrial fibrillation; HFA-PEFF, Heart Failure Association Pretest assessment, echocardiography and natriuretic peptide, functional testing and final aetiology; HFpEF, heart failure preserved left ventricular ejection fraction.

2.7% at 5 years, $p=0.026$) (online supplemental figure S3B-D). Furthermore, HFpEF by the H₂FPEF score was not associated with MACCE in the multivariate analyses (online supplemental table S3).

DISCUSSION

We evaluated the prognostic impact of the HFA-PEFF score in a large retrospective cohort of 1679 patients who underwent CA to treat AF. The main findings were: (i)

while only 7.7% of patients were initially identified as having heart failure, 32.3% were diagnosed with HFpEF based on the HFA-PEFF score and (ii) at the 5-year assessment, HFpEF by HFA-PEFF score was an independent predictor of MACCE.

In the present study, HFpEF by HFA-PEFF score was present in 32.3% of patients. HFpEF is estimated to affect approximately 50% of patients with HF and is associated with age, sex and many comorbidities, including hypertension, diabetes mellitus, chronic kidney disease, AF and multimorbidity.¹⁹ The HFA-PEFF score was proposed as a second step based on echocardiographic parameters and biomarkers. High scores are useful in diagnosing HFpEF⁶; however, the HFA-PEFF score alone provides insufficient information on the risk factors for HFpEF and exercise intolerance. On the other hand, the H₂FPEF score was established based on simple clinical characteristics and echocardiographic parameters, and high scores are useful for discriminating HFpEF from non-cardiac dyspnoea regardless of biomarker.⁵ HFpEF met only the H₂FPEF score was considered early HFpEF with obesity, hypertension, a low biomarker and a few findings indicating elevated left atrial pressure. HFpEF, according to both scores, may be distinguished between HFpEF phenotypes with different backgrounds. The results of the present study might suggest that HFA-PEFF score has a potentially superior sensitivity to identify progressive stages of HFpEF compared with the H₂FPEF score.

AF was associated with an increased risk of all-cause mortality and hospitalisation for HF and stroke, regardless of EF.²⁰ In patients with HFpEF according to LVEF and history of HF, a retrospective, multicentre study has shown that the 3-year cumulative risk for the clinical event was 11.6%, which was lower than that of patients with reduced LVEF.²¹ In the present study, the MACCE in patients with HFpEF according to HFA-PEFF score was 11.2% at 5 years, which was lower than that of previous studies,^{11 12 21} as there were fewer hospitalisations for HF. Nevertheless, HFpEF by the HFA-PEFF score was useful in predicting all-cause death, hospitalisation for HF and hospitalisation for stroke. Several studies have reported that the HFA-PEFF score is significantly inferior to the H₂FPEF score for diagnostic accuracy^{7 22}; however, both scores have been associated with clinical events in patients with HFpEF.^{23–25} There have been few studies on CA and clinical events in HFpEF using the HFA-PEFF score. CA reduces hospitalisations for HF and HF symptoms and improves diastolic function compared with medical therapy in patients with HFpEF as measured by the HFA-PEFF score.^{11 12} In the present study, the HFA-PEFF score independently predicted MACCE in patients with HFpEF undergoing CA. Importantly, HFpEF by the HFA-PEFF score showed an independent prognostic value of MACCE, while HFpEF by the H₂FPEF score was not independently associated with HF and ischaemic stroke, in contrast with the previous study.²⁶ The significant difference between the two scores is the use of biomarkers, which can be the most objective indicators of the factors

and influence prognostic evaluation.²⁷ Thus, the HFA-PEFF score may exhibit a propensity towards identifying HFpEF in more advanced stages, whereas the H₂FPEF score could potentially identify HFpEF at an earlier stage, particularly in patients without elevated BNP levels.²⁸ These findings suggest that CA for ‘early HFpEF’, which is characterised as non-HFpEF by the HFA-PEFF score and HFpEF by the H₂FPEF score, might contribute to a better prognosis through the prevention of HF and stroke.

Study limitations

This study has several limitations. First, it is a single-centre retrospective study. The results of this study may not be generalised because these patients represent a population of patients who have undergone CA. Second, the lack of information on the severity of HF symptoms and medications might have influenced the clinical events. However, the study population was not taking the sodium-glucose co-transporter-2 inhibitors, which have been reported to improve prognosis in patients with HFpEF,^{29 30} at the start of study observation. Third, the HFA-PEFF score might have been inaccurate because it did not use global longitudinal strain, which might have led to underestimation. Conversely, other factors in the HFA-PEFF score were considered accurate as echocardiographic examinations were performed by expert echocardiographers and physicians.

CONCLUSION

The results of our study suggest that the early detection of HFpEF using the HFA-PEFF score may contribute to cardiovascular and cerebrovascular risk prediction in patients with AF undergoing CA. This finding may have clinical applications for therapeutic decision-making for CA by detecting ‘early HFpEF’ regardless of HF symptoms. There is a need for further studies to test our findings in a larger population or multicentre, prospective study in the future.

Acknowledgements We would like to acknowledge the Department of Cardiovascular Medicine and the Department of Clinical Laboratory for their cooperation in obtaining the data for this study. Moreover, we thank Editage (www.editage.com) for the English-language editing of this manuscript.

Contributors TO structured and drafted the manuscript. The other authors drafted, reviewed and edited the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Consent obtained directly from patient(s).

Ethics approval The study protocol conformed to the ethical principles outlined in the Declaration of Helsinki and was approved by the Medical Research Ethics Committee at Kobe City Medical Center General Hospital (k230307). Moreover, an opt-out system was used to obtain patients’ consent for the use of their clinical data for research purposes.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those

of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Taiji Okada <http://orcid.org/0000-0002-8878-3446>

Toshiaki Toyota <http://orcid.org/0000-0002-7214-2265>

Tomohiko Taniguchi <http://orcid.org/0000-0002-4999-3432>

REFERENCES

- Vermond RA, Geelhoed B, Verweij N, *et al.* Incidence of atrial fibrillation and relationship with cardiovascular events, heart failure, and mortality: A community-based study from the Netherlands. *J Am Coll Cardiol* 2015;66:1000–7.
- Zafir B, Lund LH, Laroche C, *et al.* Prognostic implications of atrial fibrillation in heart failure with reduced, mid-range, and preserved ejection fraction: a report from 14 964 patients in the European society of cardiology heart failure long-term Registry. *Eur Heart J* 2018;39:4277–84.
- McDonagh TA, Metra M, Adamo M, *et al.* 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure: developed by the task force for the diagnosis and treatment of acute and chronic heart failure of the European society of cardiology (ESC) with the special contribution of the heart failure Association (Hfa) of the ESC. *Rev Esp Cardiol (Engl Ed)* 2022;75:523.
- Bozkurt B, Coats AJ, Tsutsui H, *et al.* Universal definition and classification of heart failure: A report of the heart failure society of America, heart failure Association of the European society of cardiology, Japanese heart failure society and writing committee of the universal definition of heart failure. *J Card Fail* 2021:00050–6.
- Reddy YNV, Carter RE, Obokata M, *et al.* A simple, evidence-based approach to help guide diagnosis of heart failure with preserved ejection fraction. *Circulation* 2018;138:861–70.
- Barandiarán Aizpurua A, Sanders-van Wijk S, Brunner-La Rocca H-P, *et al.* Validation of the HFA-PEFF score for the diagnosis of heart failure with preserved ejection fraction. *Eur J Heart Fail* 2020;22:413–21.
- Tada A, Nagai T, Omote K, *et al.* Performance of the H2Fpef and the HFA-PEFF scores for the diagnosis of heart failure with preserved ejection fraction in Japanese patients: A report from the Japanese multicenter Registry. *Int J Cardiol* 2021;342:43–8.
- Sanders-van Wijk S, Barandiarán Aizpurua A, Brunner-La Rocca H-P, *et al.* The HFA-PEFF and H2 FPEF scores largely disagree in classifying patients with suspected heart failure with preserved ejection fraction. *Eur J Heart Fail* 2021;23:838–40.
- Verbrugge FH, Reddy YNV, Sorimachi H, *et al.* Diagnostic scores predict morbidity and mortality in patients hospitalized for heart failure with preserved ejection fraction. *Eur J Heart Fail* 2021;23:954–63.
- Selvaraj S, Myhre PL, Vaduganathan M, *et al.* Application of diagnostic Algorithms for heart failure with preserved ejection fraction to the community. *JACC Heart Fail* 2020;8:640–53.
- Rattka M, Kühberger A, Pott A, *et al.* Catheter ablation for atrial fibrillation in Hfpef patients-A propensity-score-matched analysis. *J Cardiovasc Electrophysiol* 2021;32:2357–67.
- Rattka M, Pott A, Kühberger A, *et al.* Restoration of sinus rhythm by pulmonary vein isolation improves heart failure with preserved ejection fraction in atrial fibrillation patients. *Europace* 2020;22:1328–36.
- Arora S, Jaswaney R, Jani C, *et al.* Catheter ablation for atrial fibrillation in patients with concurrent heart failure. *Am J Cardiol* 2020;137:45–54.
- Packer DL, Piccini JP, Monahan KH, *et al.* Ablation versus drug therapy for atrial fibrillation in heart failure: results from the CABANA trial. *Circulation* 2021;143:1377–90.
- Otto CM, Nishimura RA, Bonow RO, *et al.* ACC/AHA guideline for the management of patients with valvular heart disease: A report of the American college of cardiology/American heart Association joint committee on clinical practice guidelines. *Circulation* 2021;143:e72–227.
- Lang RM, Badano LP, Mor-Avi V, *et al.* Recommendations for cardiac chamber Quantification by echocardiography in adults: an update from the American society of echocardiography and the European Association of cardiovascular imaging. *J Am Soc Echocardiogr* 2015;28:1–39.
- Nagueh SF, Smiseth OA, Appleton CP, *et al.* Recommendations for the evaluation of left ventricular diastolic function by echocardiography: an update from the American society of echocardiography and the European Association of cardiovascular imaging. *J Am Soc Echocardiogr* 2016;29:277–314.
- Sumida T, Tanabe K, Yagi T, *et al.* Single-beat determination of Doppler-derived aortic flow measurement in patients with atrial fibrillation. *J Am Soc Echocardiogr* 2003;16:712–5.
- Gevaert AB, Kataria R, Zannad F, *et al.* Heart failure with preserved ejection fraction: recent concepts in diagnosis, mechanisms and management. *Heart* 2022;108:1342–50.
- Son MK, Park JJ, Lim N-K, *et al.* Impact of atrial fibrillation in patients with heart failure and reduced, mid-range or preserved ejection fraction. *Heart* 2020;106:1160–8.
- Fujimoto H, Doi N, Okayama S, *et al.* Long-term prognosis of patients undergoing radiofrequency catheter ablation for atrial fibrillation: comparison between heart failure subtypes based on left ventricular ejection fraction. *Europace* 2022;24:576–86.
- Reddy YNV, Kaye DM, Handoko ML, *et al.* Diagnosis of heart failure with preserved ejection fraction among patients with unexplained Dyspnea. *JAMA Cardiol* 2022;7:891–9.
- Parcha V, Malla G, Kalra R, *et al.* Diagnostic and Prognostic implications of heart failure with preserved ejection fraction scoring systems. *ESC Heart Fail* 2021;8:2089–102.
- Hashemi D, Mende M, Trippel TD, *et al.* Evaluation of the HFA-PEFF score: results from the prospective DIAST-CHF cohort. *ESC Heart Fail* 2022;9:4120–8.
- Przewlocka-Kosmala M, Butler J, Donal E, *et al.* Prognostic value of the MAGGIC score, H2Fpef score, and HFA-PEFF algorithm in patients with Exertional Dyspnea and the incremental. *Journal of the American Society of Echocardiography* 2022;35:966–75.
- Kim M, Yu HT, Kim T-H, *et al.* Ischemic $t_{s_{ro}}$ Ke in non-gender-related Cha2Ds2-VA score 0–1 is associated with H2Fpef score among the patients with atrial fibrillation. *Front Cardiovasc Med* 2022;8:791112.
- Hendricks S, Dykun I, Balcer B, *et al.* Higher BNP/NT-Pro BNP levels stratify prognosis equally well in patients with and without heart failure: a Meta-Analysis. *ESC Heart Fail* 2022;9:3198–209.
- Shah Z, Wiley M, Sridhar AM, *et al.* Inverse correlation of venous brain natriuretic peptide levels with body mass index is due to decreased production. *Cardiology* 2017;137:159–66.
- Anker SD, Butler J, Filippatos G, *et al.* Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med* 2021;385:1451–61.
- Solomon SD, McMurray JJV, Claggett B, *et al.* Dapagliflozin in heart failure with mildly reduced or preserved ejection fraction. *N Engl J Med* 2022;387:1089–98.

Supplementary table S1. Univariable and multivariable analyses for MACCE for the sensitivity analysis

	Univariate analysis		Sensitive analysis	
	HR (95% CI)	<i>P</i>	HR (95% CI)	<i>P</i>
HFpEF by the HFA-PEFF score	2.56 (1.63 – 4.01)	< 0.001	1.65 (1.03 – 2.64)	0.038
Age	1.09 (1.05 – 1.12)	< 0.001	1.07 (1.04 – 1.10)	< 0.001
Body mass index	0.94 (0.87 – 1.00)	0.058		
Body surface area	0.15 (0.05 – 0.49)	0.002		
Chronic kidney disease \geq grade 3	1.95 (1.25 – 3.04)	0.003		
LV end-diastolic dimension	0.96 (0.91 – 1.01)	0.077		
Moderate or severe TR	2.80 (1.65 – 4.75)	< 0.001	1.83 (1.07 – 3.15)	0.028

Chronic kidney disease is defined as eGFR < 60 mL/min/1.73 m².

CI, confidence interval; *GFR*, glomerular filtration rate; *HFpEF*, heart failure with preserved left ventricular ejection fraction; *HR*, hazard ratio; *TR*, tricuspid regurgitation.

Supplementary table S2. Characteristics and outcome of the study population divided by HFpEF or non-HFpEF in the H₂FPEF score

	HFpEF	Non-HFpEF	P*
	(n = 560)	(n = 1119)	
Demographic data:			
Age, years	75 ± 8	69 ± 11	< 0.001
Sex, male	355 (63.4)	863 (77.1)	< 0.001
Height, m	1.62 ± 0.09	1.67 ± 0.09	< 0.001
Weight, kg	66.1 ± 14.2	65.3 ± 11.3	0.256
Body mass index, kg/m ²	24.9 ± 4.2	23.2 ± 2.9	< 0.001
Body surface area, m ²	1.70 ± 0.20	1.73 ± 0.18	0.003
CHADS ₂ score	2 (1 – 2)	1 (0 – 2)	< 0.001
Congestive heart failure	59 (10.5)	71 (6.3)	0.002
Hypertension	526 (93.9)	399 (35.7)	< 0.001
Age ≥ 75 years	298 (53.2)	366 (32.7)	< 0.001
Diabetes mellitus	112 (19.6)	135 (12.1)	< 0.001
Prior stroke	61 (10.9)	86 (7.7)	0.028
Coronary artery disease	60 (10.8)	82 (7.4)	0.020
Chronic kidney disease ≥ grade 3	23 (41.9)	354 (31.8)	< 0.001
Paroxysmal AF	363 (64.8)	722 (64.5)	0.904

2D echocardiographic data:

LV end-diastolic dimension, mm	46.1 ± 4.6	45.5 ± 4.5	0.010
LVEF, %	62.8 ± 4.0	62.1 ± 4.0	0.001
LV mass index, g/m ²	87.9 ± 17.9	79.6 ± 15.8	< 0.001
LA diameter, mm	40.3 ± 6.4	37.2 ± 6.2	< 0.001
LA volume index, mL/m ²	47.3 ± 14.9	41.3 ± 13.7	< 0.001
LA emptying fraction, %	31.4 ± 14.7	33.1 ± 14.7	0.028
E velocity, cm/s	83.7 ± 19.9	71.3 ± 18.4	< 0.001
Septal e', cm/s	6.8 ± 1.9	8.4 ± 2.3	< 0.001
Lateral e', cm/s	8.8 ± 2.5	10.7 ± 2.9	< 0.001
Mean E/e'	11.3 ± 2.9	7.8 ± 2.4	< 0.001
TRV, m/s (N = 1547)	2.5 ± 0.3	2.3 ± 0.3	< 0.001
Moderate or severe TR	73 (13.1)	88 (7.9)	0.001
Laboratory data:			
BNP, pg/mL (N = 826)	55.4 (33.9 – 123.1)	45.9 (20.6 – 96.2)	< 0.001
NT pro-BNP, pg/mL (N = 942)	280.0 (97.0 – 708.2)	156.2 (56.7 – 538.9)	< 0.001
Estimated GFR, mL/min/1.73 m ²	63.2 ± 14.5	66.9 ± 14.5	< 0.001
HFA-PEFF score	5 (4 – 6)	3 (2 – 4)	< 0.001

Outcome:	6 (6 – 6)	5 (4 – 5)	< 0.001
5-year MACCE			
5-year death	33 (5.9)	44 (3.9)	0.070
5-year hospitalisation for heart failure	19 (3.4)	19 (1.7)	0.028
5-year hospitalisation for stroke	5 (0.9)	6 (0.5)	0.393
5-year AF recurrence	13 (2.3)	21 (1.9)	0.542
	176 (31.4)	318 (28.5)	0.216

Data are expressed as mean \pm SD or median (interquartile range) or number (%).

**P* values refer to the difference between HFpEF and non-HFpEF.

AF, atrial fibrillation; BNP, B-type natriuretic peptide; GFR, glomerular filtration rate; HFpEF, heart failure preserved left ventricular ejection fraction; LA, left atrial; LV, left ventricular; LVEF, left ventricular ejection fraction; MACCE, major adverse cardiovascular and cerebrovascular events; NT pro-BNP, N-terminal prohormone brain natriuretic peptide; TR, tricuspid regurgitation; TRV, tricuspid regurgitation peak velocity.

Okada et al.

Supplementary table S3. Univariable and multivariable analyses for MACCE for the H₂FPEF score

	Univariate analysis		Multivariate analysis 1		Multivariate analysis 2	
	HR (95% CI)	<i>P</i>	HR (95% CI)	<i>P</i>	HR (95% CI)	<i>P</i>
HFpEF by the H ₂ FPEF score	1.53 (0.97 – 2.40)	0.067	1.08 (0.67 – 1.72)	0.759	1.04 (0.65 – 1.67)	0.872
Age	1.09 (1.05 – 1.12)	< 0.001	1.08 (1.05 – 1.12)	< 0.001	1.07 (1.04 – 1.11)	< 0.001
Male	0.73 (0.46 – 1.18)	0.199	0.99 (0.61 – 1.60)	0.952	1.04 (0.64 – 1.70)	0.882
Diabetes mellitus	0.97 (0.51 – 1.83)	0.922			0.94 (0.50 – 1.79)	0.853
Prior stroke	1.57 (0.81 – 3.04)	0.186			1.35 (0.69 – 2.63)	0.381
Chronic kidney disease ≥grade 3	1.95 (1.25 – 3.04)	0.003			1.37 (0.86 – 2.16)	0.183
Moderate or severe TR	2.80 (1.65 – 4.75)	< 0.001			1.94 (1.13 – 3.33)	0.017


Multivariate analysis model 1 includes HFpEF, age, and sex. Multivariate analysis model 2 includes model 1, hypertension, diabetes mellitus, prior stroke, chronic kidney disease defined as eGFR < 60 mL/min/1.73m², and moderate or severe TR. *CI*, confidence interval; *GFR*, glomerular filtration rate; *HFpEF*, heart failure with preserved left ventricular ejection fraction; *HR*, hazard ratio; *TR*, tricuspid regurgitation.

Okada et al.

Supplementary Figure S1

Description of the HFA-PEFF score (A), and the distribution of the HFA-PEFF score (B) in this study.

A	Functional	Morphological	Biomarker (SR)	Biomarker (AF)
Major criteria: 2 points	E' (sep) <7 cm/s or E'(lat) <10 cm/s or Mean E/e' ≥15 or TRV >2.8 m/s	LAVI >34 mL/m ² or LVMI ≥149/122 g/m ² (M/F) and RWT >0.42	BNP >80 pg/mL or NT-pro-BNP >220 pg/mL	BNP >240 pg/mL or NT-pro-BNP >660 pg/mL
Minor criteria: 1 point	Mean E/e' 9–14	LAVI 29–34 mL/m ² or LVMI >115/95 g/m ² (M/F) or RWT >0.42 or LV wall thickness ≥12 mm	BNP 35–80 pg/mL or NT-pro-BNP 125–220 pg/mL	BNP 105–240 pg/mL or NT-pro-BNP 365–660 pg/mL
HFA-PEFF score ≥5 points: HFpEF (Eur Heart J. 2019; 40: 3297–3317)				




AF, atrial fibrillation; BNP, B-type natriuretic peptide; HFpEF, heart failure preserved left ventricular ejection fraction; LAVI, left atrial volume index; LV, left ventricular; LVMI, left ventricular mass index; RWT, relative wall thickness; TRV, tricuspid regurgitation peak velocity.

Okada et al.

Supplementary Figure S2Description of the H₂FPEF score (A), and the distribution of the H₂FPEF score (B) in this study.

A	Clinical Variable	Values	Points
H2	Heavy	Body mass index >30 kg/m ²	2
	Hypertensive	Hypertension or treatment with two or more anti-hypertensive drugs	1
F	AF	Paroxysmal or Persistent	3
P	Pulmonary Hypertension	TRV >2.8 m/s	1
E	Elder	Age >60 years	1
F	Filling pressure	E/e' ratio >9	1

H₂FPEF score ≥6 points: HFpEF (Circulation. 2018; 138: 861–870)




AF, atrial fibrillation; HFpEF, heart failure preserved left ventricular ejection fraction; TRV, tricuspid regurgitation peak velocity.

Okada et al.

Supplementary Figure S3

Kaplan–Meier analyses show the cumulative rate of the primary composite endpoint (A), all-cause death (B), hospitalisation for HF (C), and hospitalisation for stroke (D) stratified by the H₂FPEF score.



HFpEF, heart failure preserved left ventricular ejection fraction.