openheart Variability of cardiovascular magnetic resonance (CMR) T1 mapping parameters in healthy volunteers during long-term follow-up

Fabian aus dem Siepen,¹ Christian Baumgärtner,¹ Matthias Müller-Hennessen,¹ Florian André,¹ Daniel Messroghli,² Marco Ochs,¹ Johannes Riffel,¹ Evangelos Giannitsis,¹ Hugo A Katus,¹ Matthias G Friedrich,¹ Sebastian J Buss¹

To cite: aus dem Siepen F, Baumgärtner C, Müller-Hennessen M, *et al.* Variability of cardiovascular magnetic resonance (CMR) T1 mapping parameters in healthy volunteers during long-term follow-up. *Open Heart* 2018;**5**:e000717. doi:10.1136/openhrt-2017-000717

FS and CB contributed equally.

Received 7 September 2017
Accepted 20 December 2017

¹Department of Cardiology, Angiology and Respiratory Medicine, University Hospital Heidelberg, Heidelberg, Germany

²Department of Internal Medicine – Cardiology, Deutsches Herzzentrum Berlin and Charité, University Medicine Berlin, Berlin, Germany

Correspondence to

Dr Fabian aus dem Siepen; fabian.siepen@med.uni-heidelberg.de

ABSTRACT

Background Myocardial T1 and extracellular volume (ECV) derived from cardiovascular MRIs are more and more widely accepted as important markers for diagnosis, risk prediction and monitoring of cardiac disease. Yet data regarding long-term stability of myocardial T1 mapping are lacking. The aim of this study was to investigate the long-term stability of native and postcontrast T1 mapping values in healthy volunteers.

Methods 18 strictly selected healthy volunteers (52±10 years, 12 men) were studied on a Philips Achieva 1.5 Tesla scanner. T1 relaxation times were measured before and 15 min after a bolus contrast injection of gadolinium diethylenetriamine penta-acetic acid (DTPA) (0.2 mmol/kg) using a single-breath-hold modified Look-Locker inversion recovery 3(3)3(3)5 sequence. ECV was calculated using native and postcontrast T1 times of myocardium and blood correcting for blood haematocrit. Exams were repeated 3.6±0.5 years later under the same conditions and using the same scan protocols.

Results Cardiac biomarkers (high-sensitivity troponin T and N terminal pro-brain natriuretic peptide) remained unchanged, as well as left ventricular mass, and global and longitudinal function. No significant change occurred regarding native T1 times (1017±24 ms vs 1015±21 ms; P=0.6), postcontrast T1 times (426±38 ms vs 413±20 ms; P=0.13) or ECV (22%±2% vs 23%±2%; P=0.3). Native T1 time and ECV appeared to be better reproducible than postcontrast T1, resulting in lower coefficients of variation (ECV: 3.5%, native T1: 1.3%, postcontrast T1: 6.4%) and smaller limits of agreement (ECV: 2%/-2%, native T1: 39 ms/-35 ms, postcontrast T1: 85 ms/-59 ms).

Conclusions During long-term follow-up, native T1 and ECV values are very robust markers, whereas postcontrast T1 results appear less stable.

INTRODUCTION

Background

Cardiovascular magnetic resonance (CMR) T1 mapping represents a quantitative imaging technique for the characterisation of the myocardium and identifies changes in myocardial tissue composition. T1 mapping therefore offers the opportunity to detect

Key questions

What is already known about this subject?

- Myocardial T1 and extracellular volume (ECV) derived from cardiovascular MRIs are more and more widely accepted as important markers for diagnosis, risk prediction and monitoring of cardiac disease. Yet data regarding long-term stability of myocardial T1 mapping are lacking.

What does this study add?

- This study investigated the long-term stability of native and postcontrast T1 mapping values in healthy volunteers.

How might this impact on clinical practice?

- The demonstrated long-term stability of T1 mapping parameters in this study might help to establish this technique in clinical routine.

and to quantify pathological changes of the myocardium in various heart diseases.¹ In T1 mapping, parametric maps are generated where the signal intensity of each pixel represents its longitudinal or spin-lattice relaxation time T1 in milliseconds, which is a time constant that describes a specific magnetic tissue property. T1 maps can be assessed before (native T1) and after (postcontrast T1) administration of a gadolinium-based extracellular contrast agent. Moreover, the extracellular volume (ECV) fraction can be estimated from a combination of native and postcontrast T1 values.

Recently, several studies could demonstrate characteristic changes in native myocardial T1, postcontrast T1 and ECV in different forms of ischaemic and non-ischaemic heart diseases, including acute² and early states of diseases.³ For instance, changes in T1 times can be caused by interstitial fibrosis,^{4,5} oedema⁶ or amyloid deposition.⁷

The capability of this technique to quantify the amount of diffuse cardiac fibrosis could be demonstrated by comparing T1 mapping parameters with histological parameters from myocardial biopsies.^{8–10} Recent studies revealed even a prognostic value for T1 mapping in patients with aortic stenosis.¹¹ Furthermore, T1 mapping might be able to detect early microvascular dysfunction in adenosine stress tests.¹²

Before a diagnostic test is implemented in clinical routine, its technical characteristics have to be investigated, in particular with regard to normal values, comparability, reproducibility and stability. Recent studies reported normal ranges for healthy subjects and data on short-term stability for 1.5 and 3 Tesla. Additionally, Piechnik *et al.*¹³ reported only small differences in native T1 in a cohort of 18 healthy volunteers over a period of 3 years using the shortened modified Look-Locker inversion recovery (MOLLI) technique.

Study aim

To date, it is unknown however if ECV values of healthy individuals change over time or remain constant. We sought to investigate the long-term stability of T1 mapping values including ECV in a cohort of well-characterised, healthy volunteers.

METHODS

Study population

Eighteen volunteers were prospectively enrolled between 2010 and 2011. Recruitment was performed using postings and newspaper advertisement, inviting apparently healthy volunteers to participate in a scientific study and following a standard protocol. Except for thyroid hormone substitution and oral contraceptives, permanent medication was a contraindication. For follow-up, these volunteers were enrolled by written invitation in 2015. All participants gave written informed consent.

Clinical examination and ECG

All subjects underwent clinical examination and 12-lead ECG at the baseline visit and at the follow-up visit. Medical history, family history, physical examination and ECG analysis were performed by two clinically experienced physicians and supervised by senior physicians.

Laboratory testing

Extensive blood analyses were performed both at the baseline visit and at the follow-up visit, including complete blood cell count, electrolytes, cardiac biomarkers, liver and kidney function tests, and thyroid hormone levels. Additionally, test subjects underwent an oral glucose tolerance test at the baseline visit. Standard assays were used for determination of troponin T (ECLIA, Roche Diagnostics) and Nterminal pro-brain natriuretic peptide (NT-proBNP) (CLIA, Siemens). Blood samples were collected on the same day as the CMR scan was performed.

Cardiovascular magnetic resonance

All scans were performed on the same Philips 1.5 Tesla MRI scanner (Achieva) with a 32-element cardiac receiver coil and vector-ECG gating. Steady-state free-precession cine sequences were applied in contiguous short-axis and in three long-axis views (two-chamber, three-chamber, four-chamber). T1 imaging was performed using a MOLLI sequence in a mid-ventricular short-axis view during breath-hold in end-expiration¹⁴ before and 15 min after administration of gadolinium diethylenetriamine penta-acetic acid /DTPA (Magnevist, 0.2 mmol/kg body weight). At baseline, an adenosine stress examination was performed to exclude significant coronary artery disease.

Contiguous short-axis stack images were used to measure cardiac volumes, left ventricular ejection fraction (LVEF) and myocardial mass in late diastole using a commercially available workstation (Philips IntelliSpace, V.7.0.1, Best, The Netherlands).

Precontrast and postcontrast T1 maps were generated and analysed using a commercial software package for cardiovascular image analysis (cvi⁴², V.5.2, Circle Cardiovascular Imaging, Calgary, Canada). Endocardial and epicardial borders were defined manually, using an offset of 15% to avoid partial-volume effects in the subendocardial and subepicardial layers. Global myocardial T1 values were calculated as a mean of all segments with respect to the segment's area. T1 values for blood were gathered by manually drawing a region of interest in the left ventricular (LV) cavity.

ECV values were created according to the following formula¹⁵:

$$ECV = (1 - HCT) \frac{\left(\frac{1}{T1_{myo_{post}}} - \frac{1}{T1_{myo_{pre}}} \right)}{\left(\frac{1}{T1_{blood_{post}}} - \frac{1}{T1_{blood_{pre}}} \right)}$$

Statistical analysis

Statistical analysis was performed using MedCalc V.16.4 statistical software (MedCalc Software, Ostend, Belgium).

The Kolmogorov-Smirnov test was used to test for normal distributions. Categorical data are expressed as percentages and continuous variables as mean±SD. For comparison of two parametric variables, the paired t-test was used. For analysis of the reproducibility of different parameters, we used Bland-Altman plots and coefficients of variation. A P value of less than 0.05 was considered statistically significant.

RESULTS

Clinical parameters and laboratory results

No study subjects reported symptoms or any other evidence of cardiac disease at baseline and after follow-up (mean interval 3.8±0.6 years). None of the subjects had medications on a regular basis. All clinical and ECG values were within the normal range at baseline as well as at the follow-up visit. There was a small yet significant increase of systolic (from 119±10 to 126±9 mm Hg;

Table 1 Clinical characteristics and ECG data

	Baseline visit	Follow-up visit	P value
Height (cm)	174±11	174±11	0.67
Weight (kg)	76±16	76±16	0.42
Body mass index (kg/m ²)	25±3	25±3	0.43
Age (years)	54±8	58±8	<0.01
Systolic blood pressure (mm Hg)	119±10	126±9	<0.001
Diastolic blood pressure (mm Hg)	73±10	83±8	<0.001
ECG parameters			
Heart rate (1/min)	67±9	62±6	<0.01
PQ (ms)	158±25	163±25	0.06
QRS (ms)	98±10	95±7	0.1
QTc (ms)	398±12	405±10	0.05

Continuous data were expressed as mean±SD.

P<0.001) and diastolic (from 73±10 to 83±8 mm Hg; P<0.001) blood pressure, PQ time, QTc time, and heart frequency between the two examinations, while all other parameters remained stable (table 1). No arrhythmias, abnormalities in conduction or signs of ischaemia were detected by ECG.

Laboratory values were within normal range at both visits, and the detailed data are provided in table 2. No significant changes in cardiac biomarkers (high-sensitivity troponin T and NT-proBNP) were observed.

CMR results

All CMR parameters of all subjects were within the reference range on both visits, with normal initial stress test results. On the follow-up visit, significant changes occurred in end-diastolic volume (from 155±31 mL to 143±28 mL, P=0.01), end-systolic volume (from 60±14 mL to 55±12 mL, P=0.01) and stroke volume (from 94±20 mL to 88±17 mL, P=0.03), whereas LV mass, LVEF and parameters of longitudinal function remained unchanged. All parameters are given in table 3.

No significant change occurred in native T1 times (baseline: 1017±24 ms vs follow-up: 1015±21 ms; P=0.6; figure 1), postcontrast T1 times (baseline: 426±38 ms vs follow-up: 413±20 ms; P=0.13; figure 2) or ECV (baseline: 22±2% vs follow-up: 23%±2%; P=0.3; figure 3). Native T1 time and ECV were better reproducible than postcontrast T1, resulting in better coefficients of variation (ECV: 3.5%, native T1: 1.3%, postcontrast T1: 6.4%). Limits of agreement were smaller in ECV and native compared with postcontrast T1 (ECV: 2%/–2%, native T1: 39 ms/–35 ms, postcontrast T1: 85 ms/–59 ms). Bland-Altman plots are given in figure 4.

Gender-wise analysis

Male subjects had a significantly higher body mass index (26±1 kg/m² vs 22±1 kg/m², P=0.02), but did not differ

Table 2 Laboratory results

	Baseline visit	Follow-up visit	P value
High-sensitivity troponin T (pg/mL)	3.9±1.3	3.9±1.1	0.99
NT-proBNP (ng/L)	68±44	71±48	0.66
GOT (U/L)	21±4	29±15	0.03
GPT (U/L)	22±9	30±36	0.39
Bilirubin (mg/dL)	0.8±0.4	0.8±0.2	0.31
Gamma-GT (U/L)	23±14	31±39	0.42
Creatinine (mg/dL)	0.9±0.1	0.8±0.1	0.02
GFR (mL/min)	96±20	96±17	0.90
Cholesterol (mg/dL)	204±23	201±33	0.44
HDL-cholesterol (mg/dL)	61±15	57±18	0.08
LDL-cholesterol (mg/dL)	127±23	123±28	0.26
Haemoglobin A1c (%)	5.5±0.2	5.3±0.2	0.76
CRP (mg/L)	2.5±1.4	2.2±0.8	0.63
Haemoglobin (g/dL)	14.2±1.2	14.3±1.3	0.45
Haematocrit (%)	42±3	41±3	<0.001
TSH (mU/L)	1.4±0.8	1.0±0.4	0.04
Uric acid (mg/dL)	5±1	5±1	0.28
Sodium (mmol/L)	141±2	141±2	0.76
Potassium (mmol/L)	4.2±0.3	4.3±0.2	0.57

Continuous data were expressed as mean±SD.

CRP, C reactive protein; GFR, glomerular filtration rate; GOT, glutamic oxaloacetic transaminase; GPT, glutamic pyruvic transaminase; Gamma-GT, Gamma-Glutamyltransferase; HDL, high-density lipoprotein; LDL, low-density lipoprotein; NT-proBNP, N-terminal pro-brain natriuretic peptide; TSH, thyroid-stimulating hormone.

significantly in any other baseline parameter. No differences in CMR parameters or T1 mapping values could be observed between genders except postcontrast T1 time at baseline visit (male: 443±8 ms, female: 388±14 ms; P=0.002). Limits of agreement were smaller in ECV and native T1 compared with postcontrast T1 in men (native T1: 24/–27 ms, postcontrast: –95/42 ms, ECV: 2%/–2%) as well as in women (native T1: 60/–51 ms, postcontrast T1: 48/–73 ms, ECV: 2%/–2%). Accordingly, coefficients of variation were better for ECV and native T1 compared with postcontrast T1 in men (ECV: 3.0%, native T1: 0.9%, postcontrast T1: 6.9%) and women (ECV: 3.9%, native T1: 1.7%, postcontrast T1: 5.9%).

DISCUSSION

In this study we could demonstrate that in a cohort of strictly selected healthy individuals, (1) native T1, postcontrast T1 and ECV did not change significantly during a long-term follow-up; (2) native T1 and ECV values showed less variation between measurements than postcontrast

Figure 1 Representative images of native T1 maps and plots of individual T1 values at baseline (t_0) and follow-up visit (t_1).

T1; and (3) no significant differences between genders could be observed.

T1 mapping is an emerging technique and recently was used in numerous single-centre studies. Significant differences between healthy and diseased subjects could be demonstrated for a wide spectrum of cardiomyopathies, including dilated cardiomyopathy (DCM),^{45 9 16} hypertrophic cardiomyopathy^{45 17–20} and cardiac amyloidosis.^{721–23} Some studies also elucidated the prognostic value of T1 mapping in a cohort of patients with DCM,^{16 24} light-chain amyloidosis,²⁵ diabetes²⁶ and patients with heart failure of different aetiologies.²⁷ An association between ECV and short-term mortality in a heterogeneous cohort could be demonstrated by Wong *et al.*²⁸ Before these research findings can be translated into clinical application, several technical aspects have to be addressed. First, various mapping sequences with specific advantages and disadvantages and—even more importantly—different normal ranges are available, and technical improvement is still ongoing. T1 results might vary between different vendors and magnetic resonance systems,²⁹ and there is a lack of multivendor/multicentre trials. Furthermore, the field is heterogeneous regarding the use and dosage of contrast agents, cut-off values, quality management, image postprocessing and reporting. Due to the novelty of the technique and the different technical challenges, no universally valid reference values are available yet. In a consensus paper edited by an expert consensus group, it was suggested that normal values should be established at each CMR site for the local set-up.³⁰ In a first multicentre, single-vendor approach, Piechnik *et al.*¹³ provided reference values for 1.5 Tesla in a large cohort of 342 healthy volunteers. Additionally, mapping values for different cardiovascular diseases were provided.³¹ Dabir *et al.*³² provided reference values for 1.5 and 3T based on 103 healthy subjects, examined in four different centres under the same conditions using a standardised mapping sequence. Normal values of 1.5T T1 mapping parameters based on smaller cohorts were also published by Messroghli *et al.*,¹⁴ Gai *et al.*,³³ Liu *et al.*,³⁴ von Knobelsdorf-Brenkenhoff *et al.*³⁵ and Nacif *et al.*³⁶ The average values in our cohort were closest to the values obtained by Nacif *et al.*, using the same sequence with the same flip angle of 35°.

Figure 2 Representative images of postcontrast T1 maps and plots of individual postcontrast T1 values at baseline (t_0) and follow-up visit (t_1).

Postcontrast T1 times depend on the dosage of contrast agents, the total volume of distribution and the time between injection of contrast medium and image acquisition. This may explain why postcontrast T1 had the highest coefficient of variation among all parameters in our cohort and therefore appeared to be less robust than native T1 or ECV. As a derived parameter, ECV appears much less dependent on technical confounders and field strength. Additionally, ECV has a lower variability between different vendors than native and post-contrast T1.²⁹ Indeed, in our study, ECV had a good

Table 3 CMR parameters

	Baseline visit	Follow-up visit	P value
LVEDD (mm)	48±4	47±3	0.3
LVESD (mm)	32±4	32±3	0.7
EDV (mL)	155±31	143±28	0.01
ESV (mL)	60±14	55±12	0.01
SV (mL)	94±20	88±17	0.03
LVEF (%)	61±4	62±3	0.4
LV mass (g)	83±24	88±26	0.06
MAPSE (mm)	12±3	12±2	0.9
TAPSE (mm)	20±3	19±3	0.5
T1 mapping parameters			
Native T1 myocardium (ms)	1017±24	1015±21	0.6
Native T1 blood (ms)	1542±87	1552±63	0.6
Postcontrast T1 myocardium (ms)	426±38	413±20	0.13
Postcontrast T1 blood (ms)	242±33	233±16	0.2
ECV (%)	22±2	23±2	0.3

Continuous data were expressed as mean±SD.

CMR, cardiovascular magnetic resonance; ECV, extracellular volume; ESV, end-systolic volume; EDV, end-diastolic volume; LV, left ventricular; LVEDD, left ventricular end-diastolic diameter; LVEF, left ventricular ejection fraction; LVESD, left ventricular end-systolic diameter; MAPSE, mitral annulus plane systolic excursion; SV, stroke volume; TAPSE, tricuspid annular plane systolic excursion.

Figure 3 Representative images of extracellular volume (ECV) maps and plots of individual ECV values at baseline (t_0) and follow-up visit (t_1).

Figure 4 Bland-Altman plots of native T1 (A), postcontrast T1 (B) and extracellular volume (ECV) (C). T_0 , baseline visit; t_1 , follow up visit.

reproducibility, with just a slightly higher coefficient of variation than native T1.

In our cohort, no significant changes could be detected between baseline and follow-up scans, but several studies reported an age dependency of T1 mapping parameters. For instance, an increase of ECV^{26 37} and a decrease of native T1 times with age¹³ were described. It remains unclear whether the observed changes might indicate a subclinical disease or the physiological ageing process. However, the small sample size in our study should be considered as a major limitation in this context.

The finding that native T1 and ECV remain constant in healthy volunteers over several years, respectively, the long-term stability of T1 mapping parameters in individuals, is an essential precondition for applications such as monitoring during therapeutic interventions or assessment of natural disease progression. Of course, our study is limited by a small case number and a lack of data for different sequences, field strengths, vendors or post-processing procedures. Nevertheless, the presented data endorse the results of previous studies regarding stability of T1 measurements.

CONCLUSION

Myocardial T1 and ECV values are stable over a long-term follow-up period in healthy subjects and thus appear suitable for longitudinal studies. While a similar reproducibility could be demonstrated for native T1 and ECV, post-contrast T1 appeared to be less robust.

Acknowledgements We thank our medical technical assistant Daniel Helm for excellent image acquisition and his valuable and sustained support.

Contributors FadS: study conception and design, drafting of manuscript. CB: acquisition of data. MM-H: analysis and interpretation of data, drafting of manuscript. FA: analysis and interpretation of data, statistics. DM: critical revision. MO: critical revision. JR: critical revision. EG: critical revision. HAK: critical revision. MGF: critical revision. SJB: analysis and interpretation of data, drafting of manuscript.

Competing interests None declared.

Ethics approval The study was conducted in accordance with the ethical principles of the Declaration of Helsinki and was approved by the local ethics committee of the University of Heidelberg, Germany (reference number: S-038/2007, Ethikkommission Heidelberg, Alte Glockengießerei 11/1, 60015 Heidelberg).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- White SK, Sado DM, Flett AS, *et al.* Characterising the myocardial interstitial space: the clinical relevance of non-invasive imaging. *Heart* 2012;98:773–9.

2. Messroghli DR, Niendorf T, Schulz-Menger J, *et al.* T1 mapping in patients with acute myocardial infarction. *J Cardiovasc Magn Reson* 2003;5:353–9.
3. Ugander M, Oki AJ, Hsu LY, *et al.* Extracellular volume imaging by magnetic resonance imaging provides insights into overt and sub-clinical myocardial pathology. *Eur Heart J* 2012;33:1268–78.
4. Puntmann VO, Voigt T, Chen Z, *et al.* Native T1 mapping in differentiation of normal myocardium from diffuse disease in hypertrophic and dilated cardiomyopathy. *JACC Cardiovasc Imaging* 2013;6:475–84.
5. Dass S, Suttie JJ, Piechnik SK, *et al.* Myocardial tissue characterization using magnetic resonance noncontrast t1 mapping in hypertrophic and dilated cardiomyopathy. *Circ Cardiovasc Imaging* 2012;5:726–33.
6. Ferreira VM, Piechnik SK, Dall'Armellina E, *et al.* Non-contrast T1-mapping detects acute myocardial edema with high diagnostic accuracy: a comparison to T2-weighted cardiovascular magnetic resonance. *J Cardiovasc Magn Reson* 2012;14:42.
7. Fontana M, Bannyersad SM, Treibel TA, *et al.* Native T1 mapping in transthyretin amyloidosis. *JACC Cardiovasc Imaging* 2014;7:157–65.
8. Sibley CT, Noureldin RA, Gai N, *et al.* T1 Mapping in cardiomyopathy at cardiac MR: comparison with endomyocardial biopsy. *Radiology* 2012;265:724–32.
9. aus dem Siepen F, Buss SJ, Messroghli D, *et al.* T1 mapping in dilated cardiomyopathy with cardiac magnetic resonance: quantification of diffuse myocardial fibrosis and comparison with endomyocardial biopsy. *Eur Heart J Cardiovasc Imaging* 2015;16:210–6.
10. Iles L, Pfluger H, Phrommintikul A, *et al.* Evaluation of diffuse myocardial fibrosis in heart failure with cardiac magnetic resonance contrast-enhanced T1 mapping. *J Am Coll Cardiol* 2008;52:1574–80.
11. Lee H, Park JB, Yoon YE, *et al.* Noncontrast Myocardial T1 Mapping by Cardiac Magnetic Resonance Predicts Outcome in Patients With Aortic Stenosis. *JACC Cardiovasc Imaging* 2017. doi: 10.1016/j.jcmg.2017.09.005. [Epub ahead of print 10 Nov 2017].
12. Levelt E, Piechnik SK, Liu A, *et al.* Adenosine stress CMR T1-mapping detects early microvascular dysfunction in patients with type 2 diabetes mellitus without obstructive coronary artery disease. *J Cardiovasc Magn Reson* 2017;19:81.
13. Piechnik SK, Ferreira VM, Lewandowski AJ, *et al.* Normal variation of magnetic resonance T1 relaxation times in the human population at 1.5 T using ShMOLLI. *J Cardiovasc Magn Reson* 2013;15:13.
14. Messroghli DR, Greiser A, Fröhlich M, *et al.* Optimization and validation of a fully-integrated pulse sequence for modified look-locker inversion-recovery (MOLLI) T1 mapping of the heart. *J Magn Reson Imaging* 2007;26:1081–6.
15. Flett AS, Hayward MP, Ashworth MT, *et al.* Equilibrium contrast cardiovascular magnetic resonance for the measurement of diffuse myocardial fibrosis: preliminary validation in humans. *Circulation* 2010;122:138–44.
16. Barison A, Grigoratos C, Todiere G, *et al.* Myocardial interstitial remodelling in non-ischaemic dilated cardiomyopathy: insights from cardiovascular magnetic resonance. *Heart Fail Rev* 2015;20:731–49.
17. Kellman P, Wilson JR, Xue H, *et al.* Extracellular volume fraction mapping in the myocardium, part 2: initial clinical experience. *J Cardiovasc Magn Reson* 2012;14:64.
18. Sado DM, Flett AS, Bannyersad SM, *et al.* Cardiovascular magnetic resonance measurement of myocardial extracellular volume in health and disease. *Heart* 2012;98:1436–41.
19. Kramer CM, Chandrashekar Y, Narula J. T1 mapping by CMR in cardiomyopathy: a noninvasive myocardial biopsy? *JACC Cardiovasc Imaging* 2013;6:532–4.
20. Mordi I, Carrick D, Bezerra H, *et al.* T1 and T2 mapping for early diagnosis of dilated non-ischaemic cardiomyopathy in middle-aged patients and differentiation from normal physiological adaptation. *Eur Heart J Cardiovasc Imaging* 2016;17:797–803.
21. Mongeon FP, Jerosch-Herold M, Coelho-Filho OR, *et al.* Quantification of extracellular matrix expansion by CMR in infiltrative heart disease. *JACC Cardiovasc Imaging* 2012;5:897–907.
22. Fontana M, Pica S, Reant P, *et al.* Prognostic Value of Late Gadolinium Enhancement Cardiovascular Magnetic Resonance in Cardiac Amyloidosis. *Circulation* 2015;132:1570–9.
23. Bannyersad SM, Sado DM, Flett AS, *et al.* Quantification of myocardial extracellular volume fraction in systemic AL amyloidosis: an equilibrium contrast cardiovascular magnetic resonance study. *Circ Cardiovasc Imaging* 2013;6:34–9.
24. Puntmann VO, Carr-White G, Jabbour A, *et al.* T1-Mapping and Outcome in Nonischemic Cardiomyopathy: All-Cause Mortality and Heart Failure. *JACC Cardiovasc Imaging* 2016;9:40–50.
25. Bannyersad SM, Fontana M, Maestrini V, *et al.* T1 mapping and survival in systemic light-chain amyloidosis. *Eur Heart J* 2015;36:244–51.
26. Wong TC, Piehler KM, Kang IA, *et al.* Myocardial extracellular volume fraction quantified by cardiovascular magnetic resonance is increased in diabetes and associated with mortality and incident heart failure admission. *Eur Heart J* 2014;35:657–64.
27. Schelbert EB, Piehler KM, Zareba KM, *et al.* Myocardial Fibrosis quantified by extracellular volume is associated with subsequent hospitalization for heart failure, death, or both across the spectrum of ejection fraction and heart failure stage. *J Am Heart Assoc* 2015;4:e002613.
28. Wong TC, Piehler K, Meier CG, *et al.* Association between extracellular matrix expansion quantified by cardiovascular magnetic resonance and short-term mortality. *Circulation* 2012;126:1206–16.
29. Raman FS, Kawel-Boehm N, Gai N, *et al.* Modified look-locker inversion recovery T1 mapping indices: assessment of accuracy and reproducibility between magnetic resonance scanners. *J Cardiovasc Magn Reson* 2013;15:64.
30. Moon JC, Messroghli DR, Kellman P, *et al.* Society for Cardiovascular Magnetic Resonance I and Cardiovascular Magnetic Resonance Working Group of the European Society of C. Myocardial T1 mapping and extracellular volume quantification: a Society for Cardiovascular Magnetic Resonance (SCMR) and CMR Working Group of the European Society of Cardiology consensus statement. *J Cardiovasc Magn Reson* 2013;15:92.
31. Liu JM, Liu A, Leal J, *et al.* Measurement of myocardial native T1 in cardiovascular diseases and norm in 1291 subjects. *J Cardiovasc Magn Reson* 2017;19:74.
32. Dabir D, Child N, Kalra A, *et al.* Reference values for healthy human myocardium using a T1 mapping methodology: results from the International T1 Multicenter cardiovascular magnetic resonance study. *J Cardiovasc Magn Reson* 2014;16:69.
33. Gai ND, Stehning C, Nacif M, *et al.* Modified Look-Locker T1 evaluation using Bloch simulations: human and phantom validation. *Magn Reson Med* 2013;69:329–36.
34. Liu S, Han J, Nacif MS, *et al.* Diffuse myocardial fibrosis evaluation using cardiac magnetic resonance T1 mapping: sample size considerations for clinical trials. *J Cardiovasc Magn Reson* 2012;14:90.
35. von Knobelsdorff-Brenkenhoff F, Prothmann M, Dieringer MA, *et al.* Myocardial T1 and T2 mapping at 3 T: reference values, influencing factors and implications. *J Cardiovasc Magn Reson* 2013;15:53.
36. Nacif MS, Turkbey EB, Gai N, *et al.* Myocardial T1 mapping with MRI: comparison of look-locker and MOLLI sequences. *J Magn Reson Imaging* 2011;34:1367–73.
37. Liu CY, Liu YC, Wu C, *et al.* Evaluation of age-related interstitial myocardial fibrosis with cardiac magnetic resonance contrast-enhanced T1 mapping: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol* 2013;62:1280–7.