

Supplementary Table 1: Independent correlates of event-free survival (via lower quartile of days alive and out-of-hospital) and all-cause re-hospitalisation including randomisation and MCI status interaction

	Lower quartile of days alive & out-of-hospital OR (95% CI)	All-cause re-hospitalisation OR (95% CI)
Socio-demographic profile		
Age (years)	1.09 (1.05 to 1.13)**	-
Risk factor profile		
noMCI-HBI	Reference	Reference*
MCI-HBI	3.38 (1.04 to 11.00)*	3.90 (1.39 to 10.90)*
noMCI-SM	2.74 (0.71 to 10.55)	2.30 (0.76 to 6.95)
MCI-SM	3.69 (1.15 to 11.85)*	5.09 (1.80 to 14.41)*
Current smoker [§]	2.42 (0.99 to 5.94)	0.37 (0.14 to 0.96)*
Co-morbidity profile		
Type 2 diabetes	2.08 (1.06 to 4.12)*	0.36 (0.15 to 0.85)*
Treatment profile		
Angiotensin II receptor blocker	-	2.45 (0.86 to 6.98)

OR = odds ratio; CI = confidence interval; MCI = mild cognitive impairment; HBI = home-based intervention; SM = standard management.
*p<0.05; **p<0.001; [§]Assessed in n=258 patients.