

openheart The introduction of refined carbohydrates in the Alaskan Inland Inuit diet may have led to an increase in dental caries, hypertension and atherosclerosis

James J DiNicolantonio, James H O'Keefe

To cite: DiNicolantonio JJ, O'Keefe JH. The introduction of refined carbohydrates in the Alaskan Inland Inuit diet may have led to an increase in dental caries, hypertension and atherosclerosis. *Open Heart* 2018;**5**:e000776. doi:10.1136/openhrt-2018-000776

Accepted 13 June 2018

Recently, DiNicolantonio found evidence that an increase in the intake of refined carbohydrates and sugar may have led to the increase in the incidence of atherosclerotic disease in the Greenland Inuit.¹ Thus, we sought to determine if a similar event occurred in the Alaskan Inuit, and in particular, the Alaskan Inland Inuit.

The traditional Alaskan Inuit lived a semi-nomadic life hunting and catching fish, marine mammals and birds. However, beginning around 1920, a group of Alaskan Inuit slowly developed a permanent settlement in the mountains coming to live in Anaktuvuk Pass in 1950.¹ In 1953, a post office was established as well as a 'white trader' store. Bang and Kristoffersen of the Gade Institute from the Department of Pathology and School of Dentistry in Norway performed two studies from two different time periods on the diet and dental health of the Alaskan Inland Inuit. They analysed the diet in 1955–1957 by 'weighing of all the food eaten by each person for two consecutive days once a month during a 2-year period. At that time, samples of the food were analysed in order to determine the content of fat, protein and carbohydrate, as well as the total caloric intake of each individual'.¹

In 1965, the dietary intake was conducted by the 'interview method' plus personal observations when visiting all of the families during meal times. Bang and Kristoffersen '...contacted all of the Eskimos involved in the study questioning each individual in detail about what food and roughly how much of each food item he consumed in the course of the year'.¹ The authors noted a dramatic increase in carbohydrate intake of 'nearly 50%' and a decrease in the intake of protein 'by about 50%' from 1955 to 1957

to 1965.¹ This rise in the intake of carbohydrate was paralleled by an 'almost 90% increase' in the sum of decayed, missing and filled permanent teeth for primary teeth (from 3.0 to 5.6) and a fourfold increase in those >6 years old, 'the percentage of caries-free persons had decreased from 74.5% to zero in 8 years...While 50% of the children in 1955–1957 had caries-free primary teeth all the children had decayed teeth in 1965... The most dramatic change was observed in individuals 30 years of age or older. In this previously caries-free group, all subjects had developed caries in 1965'.¹ Thus, the Alaskan Inland Inuit that had subsided on a diet virtually devoid of carbohydrate for most of their life who did not have any dental caries between 1955 and 1957 had all developed dental caries by 1965; in the interim, the intake of carbohydrate had increased by 50%.¹

Bang and Kristoffersen elaborated on how the Alaskan Inland Inuit diet changed, 'While previously all able men in the village frequently were out hunting, trapping and fishing, only a couple of the young men were still actively engaged in such activities in 1965. Manufacture of souvenirs had proved more profitable and the income financed the purchase of refined foods from the local stores. Consequently, *only some 20% of the food intake was made up of native foods*, mainly caribou meat. Hunting was now mostly limited to the short periods when the caribou came close to the village. In 1955–1957, the percentage of calories obtained from protein and fat was high while the percentage derived from carbohydrate was low. In 1965, the intake of protein was about 50% lower (carbohydrate intake was 50% higher'.¹ These details highlight the fact that these Eskimos were no longer



Saint Lukes Mid America Heart Institute, Kansas City, Missouri, USA

Correspondence to
Dr James J DiNicolantonio;
jjdnicol@gmail.com

subsisting on their native diet but instead a diet much higher in refined carbohydrates.

Importantly, the Anaktuvuk Pass Inuit of 1965 were 'living under the same environmental conditions as far as geographical location, climate, housing and clothing is concerned'¹ compared with those from 1955 to 1957. Thus, the most dramatic environmental change in these Alaskan Inland Inuit during the 8-year study period was the change in their diet. The dental examinations in each time period were carried out by the same investigators using a similar method under identical conditions

and it was said by these authors that an overestimation in the prevalence of dental caries in 1965 was highly unlikely as around half of the subjects were radiographed for dental caries in 1955–1957, whereas radiographs were not used in 1965. Thus, among the Alaskan Inland Inuit of Anaktuvuk Pass, an increased intake of refined carbohydrate of at least 50%, and in some estimates as much as 200%, was closely linked to a tremendous rise in dental caries.¹

'The Eskimo race is generally believed to have a low incidence of cardiovascular disease (Wilber and Levine,

Table 1 Effect of a western diet on the health of the Alaskan Inland Inuit

	Alaskan Inland Inuit 1955–1957¹ Studied by Bang and Kristoffersen	Alaskan Inland Inuit 1965¹ Studied Bang and Kristoffersen
Diet	Fat: 41% Protein: 33% Carbohydrate: 26% ($\leq 13\%^*$) 'In 1955–1957, the percentage of calories obtained from protein and fat was high while the percentage derived from carbohydrate was low'. ¹ †Importantly, the 26% carbohydrate estimate accounted for glycogen, which was not known at the time rapidly degrades to lactate on death of the animal. Thus, the carbohydrate intake was likely $<13\%$. '... they previously lived a seminomadic life stopping at places where the hunting and trapping proved best. In the summer they would migrate to the coast to fish and hunt marine mammals and birds. Within the last two to three decades, however, they have stayed permanently in the mountains'. ¹	Fat: 40% Protein: 15% Carbohydrate: 45% 'The percentage of total calories obtained from protein had <i>decreased</i> by about 50% and that from carbohydrate <i>increased</i> by nearly 50%'. ¹ 'While previously all able men in the village frequently were out hunting, trapping and fishing, only a couple of the young men were still actively engaged in such activities in 1965... the income financed the purchase of refined foods from the local stores... only some 20% of the food intake was made up of native foods, mainly caribou meat. Hunting was now mostly limited to the short periods when the caribou came close to the village'. ¹
Dental health	Prior to 1960 'few or no dental problems' in the "Eskimos living on their native food" ¹ 1955–1957 '50% of the children in 1955–1957 had caries-free primary teeth, all the children had decayed teeth in 1965... The most dramatic change was observed in individuals ≥ 30 years of age. In this previously caries-free group, all subjects had developed caries in 1965'. ¹	1965 'The change in diet was accompanied by a drastic increase in the prevalence of dental caries'. ¹ The average number of affected teeth for primary teeth showed an 'Almost 90% increase' in the sum of decayed, missing and filled permanent teeth For those '... over 6 years of age exhibited a <i>four-fold increase</i> '. ¹ 'The most dramatic change was observed in individuals 30 years of age or older. In this previously caries-free group, all subjects had developed caries in the course of 8 years'. ¹
	Alaskan Inuit (Alaskan Inland Inuit, published 1955)² Studied by Rodahl	Westernised populations
Diet	'Eskimos included in this study was about 2700 calories; the fat consumption was 105 g...' ²	Diet of USA in 1955
Blood pressure	'... no systolic blood pressure higher than 162 mm Hg was ever recorded in our 'normal' Eskimo subjects'. ² In a series of 117 Eskimo patients, only one (0.85%) of the patients had systolic blood pressure >145 mm Hg (a 60-year-old woman having a blood pressure of 200/80 mm Hg). ² 'It may be noted that Alexander (1949) found <i>hypertension to be practically non-existent among Aleuts</i> '. ² '... systolic and diastolic blood pressures were lower in Eskimos than in Whites of corresponding age'. ² '80% of the recorded systolic blood pressures were below 116 mm Hg'. ²	8.6% prevalence of hypertension in the USA (data from Dahl 1954–1956). ⁵ 11%–13% prevalence of hypertension in Chicago in 1939, which reached 25% by 1975. ⁶
Atherosclerosis and cardiovascular disease	'None of the 16 Eskimos analysed here showed any evidence of arteriosclerosis by clinical or roentgenological (X-ray) examination'. ² 'Of the entire material, one Eskimo showed calcium deposits in the arteries'. ² While it is hard to know from the paper whether this one Eskimo was out of 84 or the 9 natives above 47 to be cautious have used $n=9$, which indicates that calcium deposits in the arteries of the Alaskan Inland Eskimo (>47 years old) was just 11%, whereas prevalence of arterial calcifications is around 50% in other more westernised countries for those of similar age 40 and 49. ³ This suggests that Alaskan Inland Eskimo had an approximate 4.5 fold lower rate of arterial calcification versus other more westernised populations. In one 60-year-old Eskimo woman with a blood pressure of 200/80 mm Hg, there was slight enlargement of the left ventricle of the heart'. ² 'Dr Paul Hagglund, who has operated on a large number of Eskimos in Alaska during the last 15 years, has never seen arteriosclerosis or atherosclerosis in Eskimos (personal communication). He had the occasion to perform an autopsy on one female and two male Eskimos, aged 60–65 years, and found no arteriosclerosis'. ² 'Dr Earl Albrecht, Territory Commissioner of Health, states that arteriosclerosis is rare in Eskimos, based on clinical evidence (personal communications)'. ² '... his (Alexander 1949) electrocardiographic and clinical examination of 296 Aleuts, including 23 above the age of 60, revealed <i>almost no cardiovascular disease</i> '. ² 'Cardiovascular disease was extremely rare among the large number of Eskimo patients examined by the author (Rodahl) during a 2-year period in Alaska'. ²	71%–82% ⁷ (cases with fibrous atherosclerotic plaques) 58%–96% (cases with fibrous atherosclerotic plaques, all 14 countries studied). <i>Description:</i> 71%–82% is from New Orleans (Blacks/Caucasians, respectively), in 45–54 years old, 1960–1965. These data are from the International Atherosclerosis Project: 1960–1965 autopsy study of 23 207 sets of coronary arteries and aorta from 14 countries (table 10) in men who died of accidents, cancer, infection and miscellaneous causes 'Calcium is present in 50% of individuals aged 40 to 49% and 80% of individuals aged 60 to 69'. A higher amount of calcification suggests more advanced atherosclerotic lesions in other more westernised populations versus the Alaskan Inland Inuit 77.3% ⁸ (atherosclerotic coronary involvement). Young Korean War vets, 1953 autopsy study.

†Eliminating glycogen, which would have been rapidly depleted from the animal on death would equate to a true carbohydrate intake of $\leq 13\%$.

1950), although their diet is apparently high in fat and cholesterol'.²

Apparently, a diet high in fat and cholesterol did not seem to have negative cardiovascular consequences for the Alaskan Inuit. Rodahl published a scientific paper in 1955 on the Alaskan Inland Eskimo indicating a daily caloric intake of 2700 calories, 37% of which was fat (105 g of fat per day).² Another larger survey found that '... the average daily fat consumption in Alaskan Eskimos was 139 g (40% of the calories)'.² And the mean cholesterol intake was 340 mg daily (varying from 150 up to 700 mg per day).² These estimates were considered underestimates because the cholesterol content of certain foods was unknown at the time. Rodahl wrote, 'It may also be noted that the cholesterol intake varies greatly from one Eskimo group to another, depending on the different dietary habits. Thus it was observed that among the inland Eskimos, the Nunarmiuts at Anaktuvuk Pass, some of the men consumed as much as ≥ 70 g of boiled brain from mountain sheep in a single evening meal yielding almost 600 mg cholesterol from this food item alone'.²

Rodahl went on to write, 'In a survey of 104 Alaskan Eskimos the author found that both the systolic and diastolic blood pressures were lower in Eskimos than in Whites of corresponding age. Eighty per cent of the recorded systolic blood pressures were < 116 mm Hg and no systolic blood pressure > 162 mm Hg was ever recorded in our 'normal' Eskimo subjects. In a series of 117 Eskimo patients, only one of the patients had systolic blood pressure > 145 mm Hg'. Thus, the rate of hypertension in Alaskan Inuit appears to be virtually non-existent (1/117 being around 0.85%) and about one-tenth the rate compared with that in the USA at the time.

Rodahl also noted that, 'None of the 16 Eskimos analysed here showed any evidence of arteriosclerosis by clinical or roentgenological (X-ray) examination, and cardiovascular disease was extremely rare among the large number of Eskimo patients examined by the author during a 2-year period in Alaska'.²

Based on chest X-rays, Rodahl also noted, 'Of the entire material, one Eskimo showed calcium deposits in the arteries'.² While it is hard to know based on the paper whether this was out of 84 Inuit or the 9 natives who were older than the age of 47 years, to err on the side of caution using the 9 natives, indicates that calcium deposits in the arteries of the Alaskan Inland Inuit (> 47 years old) were at most 11%. However, the prevalence of arterial calcifications in more westernised countries for those of similar age (40–49) is approximately 50%.³ This suggests that Alaskan Inuit had around one-fifth the prevalence of arterial calcifications. As calcium deposits in the arteries generally indicate more advanced atherosclerotic lesions, and since the Inuit apparently had a lower

rate of calcifications, this may explain the lower rate of clinical cardiovascular disease based on autopsy. Indeed, it has been suggested that plaque rupture occurs at the interface between calcified and non-calcified sections of atherosclerotic plaque.³ Thus, it can be argued that a low rate of arterial calcifications in the Alaskan Inuit may have offered them protection from plaque rupture. Table 1 summarises the effects of a western diet on the health of the Alaskan Inland Inuit.

CONCLUSION

There was an approximate 50% increase in the intake of refined carbohydrates in the Alaskan Inland Inuit from 1955 to 1957 to 1965. This may have led to the dramatic rise in dental caries and subsequent increase in atherosclerosis and coronary artery disease.^{1,4} The Alaskan Inuit eating a traditional low-carbohydrate, high-fat/protein diet had a much lower incidence of atherosclerosis, hypertension and dental caries versus more westernised populations. The health decline of the Alaskan Inland Inuit may have been driven by the rise in the intake of refined carbohydrates and sugar.

Contributors JJD: wrote the initial manuscript and performed the literature search. JHO'K: provided edits to the initial manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests JJD is the author of The Salt Fix and operates the website thesaltfix.com.

Patient consent Not required.

Provenance and peer review Commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Bang G, Kristoffersen T. Dental caries and diet in an Alaskan Eskimo population. *Scand J Dent Res* 1972;80:440–4.
- 2 Rodahl K. Diet and cardiovascular disease in the Eskimos. *Trans Am Coll Cardiol* 1955;4:192–7.
- 3 Wexler L, Brundage B, Crouse J, et al. Coronary artery calcification: pathophysiology, epidemiology, imaging methods, and clinical implications. A statement for health professionals from the American Heart Association. Writing Group. *Circulation* 1996;94:1175–92.
- 4 Fodor JG, Helis E, Yazdekhasi N, et al. "Fishing" for the origins of the "Eskimos and heart disease" story: facts or wishful thinking? *Can J Cardiol* 2014;30:864–8.
- 5 Dahl LK. Possible role of salt intake in the development of essential hypertension. 1960. *Int J Epidemiol* 2005;34:967–72. discussion 72–4, 75–8.
- 6 Johnson RJ, Segal MS, Sautin Y, et al. Potential role of sugar (fructose) in the epidemic of hypertension, obesity and the metabolic syndrome, diabetes, kidney disease, and cardiovascular disease. *Am J Clin Nutr* 2007;86:899–906.
- 7 Tejada C, Strong JP, Montenegro MR, et al. Distribution of coronary and aortic atherosclerosis by geographic location, race, and sex. *Lab Invest* 1968;18:509–26.
- 8 Joseph A, Ackerman D, Talley JD, et al. Manifestations of coronary atherosclerosis in young trauma victims--an autopsy study. *J Am Coll Cardiol* 1993;22:459–67.