Correction: Role of cardiovascular imaging for the diagnosis and prognosis of cardiac amyloidosis

Agha AM, Parwani P, Guha A, *et al.* Role of cardiovascular imaging for the diagnosis and prognosis of cardiac amyloidosis. *Open Heart* 2018;5:e000881. doi:10.1136/openhrt-2018-000881

This article has been corrected since it first published. The authors want to inform the readers on the following two changes.

The second paragraph under the sub-heading ‘Late gadolinium enhancement’ of ‘Cardiovascular magnetic resonance’, should read as:

Although a very useful technique, a challenge with LGE is to choose an appropriate inversion time (TI) value. This TI value is a baseline where the myocardium is black or ‘nulled’. Incorrect determination of this null point may mask evidence of amyloidosis.²⁷ A technique called ‘phase-sensitive inversion recovery’ allows for the automated determination of an ideal TI time and may prevent user error from incorrectly masking amyloidosis on CMR²⁷ (see figure 4).

The third paragraph under the sub-heading ‘T1 mapping’ of ‘Cardiovascular magnetic resonance’, should read as:

A pre-contrast T1 time of greater than 1044 ms has been associated with a poor prognosis in AL amyloidosis.³¹ A similar cut-off of greater than 1077 ms has been associated with worse prognosis for ATTR amyloidosis, but not particularly prognostic when separated by familial and wild-type ATTR.³²

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Open Heart 2018;5. doi:10.1136/openhrt-2018-000881corr1