openheart Clinical and cost-effectiveness of the Ross procedure versus conventional aortic valve replacement in young adults

Howard Thom,¹ Alexandru Ciprian Visan,^{2,3} Edna Keeney,¹ Dan Mihai Dorobantu,^{4,5} Daniel Fudulu,² Mansour T A Sharabiani,⁶ Jeff Round,⁷ Serban Constantin Stoica²

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/openhrt-2019-001047>).

To cite: Thom H, Visan AC, Keeney E, *et al.* Clinical and cost-effectiveness of the Ross procedure versus conventional aortic valve replacement in young adults. *Open Heart* 2019;**6**:e001047. doi:10.1136/openhrt-2019-001047

Received 10 March 2019

Revised 8 April 2019

Accepted 14 April 2019

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

¹Bristol Medical School: Population Health Sciences, University of Bristol, Bristol, UK
²Cardiothoracic Surgery, Bristol Heart Institute, Bristol, UK
³Department of Cardiothoracic Surgery, University Hospital Southampton, Southampton, UK
⁴Cardiac Surgery, Bristol Royal Hospital for Children, Bristol, UK
⁵Cardiology, Institutul de Urgenta pentru Boli Cardiovasculare Prof Dr C C Iliescu, Bucuresti, Romania
⁶National Heart and Lung Institute, Imperial College London, London, UK
⁷Institute of Health Economics, Edmonton, Alberta, Canada

Correspondence to

Dr Howard Thom; howard.thom@bristol.ac.uk

ABSTRACT

Objectives In young and middle-aged adults, there are three current options for aortic valve replacement (AVR), namely mechanical AVR (mechAVR), tissue AVR (biological AVR) and the Ross operation, with no clear guidance on the best option. We aim to compare the clinical effectiveness and cost-effectiveness of the Ross procedure with conventional AVR in young and middle-aged adults.

Methods This is a systematic literature review and meta-analysis of AVR options. Markov multistate model was adopted to compare cost-effectiveness. Lifetime costs, quality-adjusted life years (QALYs), net monetary benefit (NMB), population expected value of perfect information (EVPI) and expected value of partial perfect information were estimated.

Results We identified 48 cohorts with a total number of 12 975 patients (mean age 44.5 years, mean follow-up 7.1 years). Mortality, bleeding and thromboembolic events over the follow-up period were lowest after the Ross operation, compared with mechAVR and biological AVR ($p < 0.001$). Aortic reoperation rates were lower after Ross compared with biological AVR, but slightly higher when compared with mechAVR ($p < 0.001$). At a willingness-to-pay threshold of £20 000 per QALY, the Ross procedure is more cost-effective compared with conventional AVR, with a lifetime incremental NMB of £60 952 (£30 236 to £79 464). Incremental costs were £12 323 (£6108 to £15 972) and incremental QALYs 3.66 (1.81 to 4.76). The population EVPI indicates that a trial costing up to £2.03 million could be cost-effective.

At a willingness-to-pay threshold of £20 000 per QALY, the Ross procedure is more cost-effective compared with conventional AVR, with a lifetime incremental NMB of £60 952 (£30 236 to £79 464). Incremental costs were £12 323 (£6108 to £15 972) and incremental QALYs 3.66 (1.81 to 4.76). The population EVPI indicates that a trial costing up to £2.03 million could be cost-effective.

Conclusions In young and middle-aged adults with aortic valve disease, the Ross procedure may confer greater quality of life and be more cost-effective than conventional AVR. A

Key questions

What is already known about this subject?

► The Ross operation has recently shown favourable outcomes compared with other aortic valve replacement options in young adults.

What does this study add?

► In this study we have found that the Ross operation has better outcomes, such as mortality, freedom from reintervention or freedom from embolic events, and is more economically efficient in the long term compared with conventional aortic valve replacement in the young.
 ► Additionally, we propose that a comparative clinical trial with a budget of £2 million would be warranted and cost-effective.

How might this impact on clinical practice?

► Young patients who undergo conventional aortic valve replacement, such as with mechanical or biological prosthesis, face either lifelong anticoagulation and embolic risks, or rapid valve degeneration and reinterventions, respectively.
 ► The Ross procedure is re-emerging as a better option in this age group.
 ► Lower overall costs reflect these better outcomes, but available data are only retrospective.
 ► A randomised clinical trial, shown in this study to be economically efficient, is planned to directly compare aortic valve replacement options in non-elderly patients.

high-quality randomised trial could be warranted and cost-effective.

INTRODUCTION

The standard treatment for aortic valve (AoV) disease is aortic valve replacement (AVR) when repair is not possible.¹ Options include mechanical (mechAVR) and biological (biological AVR) prostheses, or conventional AVR (cAVR) and the Ross procedure (pulmonary autograft), and rarely homo-graft valves. Bioprostheses are durable in

older patients,² with the survival advantage of mechAVR over biological AVR disappearing around the age of 60.³ However, decision making in young and middle-aged adults is challenging, with a more marked impact of cAVR on survival in younger patients,⁴ and with biological AVR and mechAVR being associated with reduced life expectancy.⁵ Furthermore, there are issues with biological valve degeneration and mechanical valve need for anticoagulation, which are more important in young patients.

Critics of the Ross procedure point out that it creates a 'two-valve disease'. However, multiple studies show very good early results and better long-term survival, comparable with that of the general population.^{6–8} The Ross procedure affords superior haemodynamics in the autograft⁹ and typically uses a homograft in the lower pressure pulmonary circulation. Despite better outcomes, the Ross procedure is underused, partly due to technical complexity. However, it can be performed safely in congenital centres and hospitals where complex aortic root operations are routine.¹⁰

Cost-effectiveness analysis compares the costs and effects of treatments, usually measured as quality-adjusted life years (QALYs), on a monetary scale and thus aids reimbursement decision making by healthcare payers.¹¹ To date there have been no cost-effectiveness analyses evaluating the Ross procedure as an option for aortic stenosis (AS) in young and middle-aged adults. Previous economic evaluations have compared cAVR and transcatheter aortic valve implantation (TAVI) with each other and with medical management.^{12–14} The results have been variable with respect to the cost-effectiveness of TAVI compared with cAVR in high-risk patients, both being cost-effective compared with medical management.^{13–15} Value of information analysis is a method to use the outputs of cost-effectiveness analyses to prioritise and design research.¹¹ Only one publication, comparing TAVI and medical management, has applied value of information analysis to AS.¹⁶

Using meta-analysis, we compared the clinical effectiveness of the Ross procedure with biological AVR and mechAVR for the treatment of young and middle-aged adults with AS. We then used these results to inform a comprehensive cost-effectiveness comparison of these procedures in England and Wales from a National Health Service (NHS) perspective. Another objective is to estimate the value of information of a randomised controlled trial comparing cAVR with the Ross procedure.

METHODS

Our model structure and inputs were informed by a review of published cost-effectiveness models evaluating cAVR and other AoV procedures. Detailed methods and results are presented in online supplementary material A. In brief, we identified 12 papers published between 2012 and 2016 comparing cAVR, TAVI and medical management. As these models did not include the Ross procedure, we adapted their structures to capture important

features specific to Ross. Input parameters, not identified in the clinical or economic systematic reviews, were informed by targeted literature searches.

Model structure and inputs

Detailed description of the model structure and the evidence base for its inputs is provided in online supplementary material B. Briefly, we adopted a discrete-time Markov multistate model, a common choice in cost-effectiveness analysis¹⁷ and in aortic stenosis.^{12–14} The model structure is illustrated and explained in online supplementary material figures S1 and S2. Patients enter the model following an initial Ross or cAVR procedure, and their states and events determine costs and QALYs per cycle. We adopt an NHS costing perspective and lifetime time horizon with a 1-year cycle length, which is consistent with previously published models.^{18–20}

We estimated the probabilities of events and transitions using a systematic literature review and meta-analysis, detailed in online supplementary material C. We searched MEDLINE, EMBASE and Cochrane Library for studies published between January 1990 and January 2016 in adults aged 18–65. A Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow chart is included in online supplementary figures S3. Data from 41 observational studies and 2 clinical trials were pooled using a random-effects meta-analysis model (online supplementary figures S4–S12 and table S2).

Several sources were used to estimate the state occupancy and event costs and utilities (online supplementary table S1), including evidence from the extensive literature on stroke prevention in atrial fibrillation.^{21–24} Costs of the cAVR and Ross procedure, and of other one-off events, were estimated according to the 2016/2017 NHS tariff. Cost estimates were assumed fixed as they represent agreed prices paid to hospitals for activity. The healthy state utility followed that of Doble *et al*,¹³ while utilities for other states and disutilities for events were adapted from the atrial fibrillation literature.^{24,25} We averaged mechAVR and biological AVR probabilities using an estimated 0.23 proportion of cAVR that is biological, with further details in online supplementary material B (table S2).

Cost-effectiveness outputs

We estimated lifetime costs and QALYs for patients undergoing cAVR and Ross procedure, as well as the net monetary benefit (NMB) and incremental NMB (INMB). The NMB is a summary of both costs and QALYs on the monetary scale; it is calculated by subtracting costs from monetary benefit, given by multiplying QALYs by a willingness-to-pay threshold, commonly assumed to be £20 000 for the UK NHS,²⁶ but often up to \$100 000 in the USA.²⁷ We performed probabilistic sensitivity analysis (PSA) to simulate 10 000 lifetime costs, QALYs, NMBs and INMBs. Means and 95% Bayesian credible intervals (CrI) were estimated for all quantities.¹¹ We generated a cost-effectiveness acceptability curve (CEAC), which plots

the probability of Ross procedure or cAVR being most cost-effective for a range of willingness-to-pay thresholds.²⁶

Value of information analysis

We estimated the per-person expected value of perfect information (EVPI), which is the expected improvement in decision making, or value, from removing all uncertainty in all parameters.¹¹ We used multilevel Monte Carlo to estimate per-person expected value of partial perfect information (EVPPI), the expected value of removing all uncertainty in only a subset of parameters.²⁸ We focused on sets of parameters likely to be informed by studies comparing cAVR with the Ross procedure. We scaled to per-population EVPI and EVPPI using a 3.5% per annum discounted 25-year technology horizon population size of 16 376 (details in online supplementary material B).

Deterministic sensitivity analyses

We conducted a one-way sensitivity analysis comparing Ross procedure with cohorts treated solely with either biological AVR or mechAVR, rather than the mixture of our base case. Also, as some costs and utility parameters were fixed, these would not be included in our primary sensitivity analyses of PSA and EVPI. These 15 parameters are listed in online supplementary material B. We conducted one-way sensitivity analyses where these parameters were set to 50% and 150% of their values.

RESULTS

Epidemiological meta-analysis results

Estimated probabilities are summarised in [table 1](#), with forest plots detailed in online supplementary material figures S4-S12. Mortality over the follow-up period was lower in Ross patients when compared with mechAVR and biological AVR (0.54%/year vs 1.38%/year and 2.54%/year, respectively; $p<0.001$). There was no evidence of difference in early mortality between groups. The Ross procedure was associated with lower

bleeding rates compared with mechAVR and biological AVR (0.11%/year vs 0.69%/year and 0.31%/year, respectively; $p<0.001$). The number of thromboembolic events was also lower following the Ross procedure in comparison with mechAVR and biological AVR (0.26%/year vs 0.86%/year and 0.57%/year, respectively; $p<0.001$). Reoperation rates on the AoV were reduced in Ross patients when compared with biological AVR, but slightly higher when compared with mechAVR (0.54%/year vs 1.28%/year and 0.37%/year, respectively; $p<0.001$). No significant differences were found in terms of infective endocarditis.

Cost-effectiveness results

Estimated means and 95% CrI from the cost-effectiveness analysis are summarised in [table 2](#). The expected lifetime incremental net benefit of £60 952 (£30 236 to £79 464) and its narrow CrI which excludes zero suggest there is strong evidence that the Ross procedure is more cost-effective than cAVR at a willingness-to-pay threshold of £20 000. These findings were repeated at a willingness-to-pay threshold of £50 000. There is also strong evidence that the Ross procedure is associated with higher lifetime QALYs, having an incremental QALY of 3.66 (1.81 to 4.76), and thus superior effectiveness to cAVR. Although the Ross procedure has higher lifetime costs than cAVR, with incremental costs of £12 323 (£6108 to £15 972), this is outweighed by the larger benefit in QALYs.

The limited uncertainty in our decision is illustrated by the CEAC ([figure 1](#)), where it appears that beyond a very low willingness-to-pay threshold (approximately £3000), there is almost 100% probability that the Ross procedure is the most cost-effective treatment option. The Ross procedure would be most cost-effective for all willingness-to-pay thresholds above £3000 per QALY, including £20 000, £50 000 and £100 000.

The deterministic sensitivity analyses showed no impact on the results or conclusions of varying the 15 parameters. The Ross procedure remained cost-effective at a

Table 1 Comparison of outcomes between the Ross operation, mechAVR and biological AVR: results from the meta-analysis

	Cohorts	Ross	mechAVR	biological AVR	P value
Early mortality (%)	46	0.24 (0.17 to 0.34)	0.3 (0.19 to 0.45)	0.28 (0.16 to 0.48)	0.75
Total mortality (%/year)	47	0.54 (0.45 to 0.64)	1.38 (1.11 to 1.74)	2.54 (1.8 to 3.59)	<0.001
AoV reoperation (%/year)	47	0.54 (0.42 to 0.69)	0.37 (0.27 to 0.5)	1.28 (0.88 to 1.85)	<0.001
Bleeding (%/year)	44	0.11 (0.06 to 0.20)	0.69 (0.5 to 0.96)	0.31 (0.14 to 0.69)	<0.001
Thromboembolic events (%/year)	44	0.26 (0.19 to 0.37)	0.86 (0.62 to 1.18)	0.57 (0.39 to 0.84)	<0.001
AoV reoperations for IE (%/year)	44	0.16 (0.12 to 0.2)	0.17 (0.1 to 0.27)	0.26 (0.18 to 0.39)	0.08
Conservatively treated IE (%/year)	46	0.21 (0.15 to 0.3)	0.27 (0.18 to 0.42)	0.29 (0.1 to 0.87)	0.64
Total IE (%/year)	39	0.34 (0.27 to 0.42)	0.44 (0.3 to 0.64)	0.63 (0.32 to 1.22)	0.14
RVOT reinterventions (%/year)	22	0.43 (0.34 to 0.54)			

Data are from random-effects models. Values in brackets are 95% CI.

AoV, aortic valve; IE, infective endocarditis; RVOT, right ventricular outflow tract; biological AVR, tissue aortic valve replacement; mechAVR, mechanical aortic valve replacement.

Table 2 Comparison of cost-effectiveness between the Ross operation and cAVR

	cAVR	Ross	Incremental values Ross vs AVR
Costs	£33 812 (£31 028 to £39 845)	£46 135 (£44 267 to £48 263)	£12 323 (£6108 to £15 972)
QALYs	11.5 (10.4 to 13.3)	15.2 (14.4 to 15.9)	3.66 (1.81 to 4.76)
NMB at £20 000	£196 385 (£177 231 to £227 366)	£257 337 (£241 933 to £270 844)	£60 952 (£30 236 to £79 464)
NMB at £50 000	£541 679 (£490 382 to £627 964)	£712 545 (£672 962 to £748 287)	£170 866 (£84 507 to £222 263)

Values represent means and CI in brackets.

cAVR, conventional aortic valve replacement; NMB, net monetary benefit; QALYs, quality-adjusted life year.

Figure 1 Cost-effectiveness acceptability curve.

willingness-to-pay threshold below £6000 for all sensitivities. Sensitivity analysis found the Ross procedure to be superior to biological AVR and mechAVR alone; this sensitivity also indicated that mechAVR was superior on costs and QALYs to biological AVR, in line with existing literature.²⁹

Value of information results

The estimated per-person expected value of information (EVPI) at willingness-to-pay £20 000 was £123.70 and 25-year population EVPI of £2.03 million. This suggests that the proposed trial design costing up to £2.03 million could be cost-effective. The EVPI results suggest that although studies on the utilities, poststroke disability costs and events informed by the meta-analysis could offer value of up to £425 000, £342 000 and £307 000, a value of up to £1.90 million could come from studying only the early event rates of stroke and bleed immediately following cAVR or Ross procedure. A trial comparing Ross and cAVR and informing the meta-analysis, early events and other epidemiological parameters, but not

costs or utilities, could approach the maximum value of £2.03 million for decision making. Value of information results are detailed in [table 3](#).

DISCUSSION

The results of our meta-analysis show lower mortality and fewer thromboembolic events for the Ross procedure versus cAVR. Rates of surgical reoperations on the AoV were lower than for biological AVR but higher than in mechAVR. Our cost-effectiveness analysis found that the Ross procedure would be the most cost-effective option for all willingness-to-pay thresholds above £3000 per QALY, due to the greater incremental effectiveness, despite the higher lifetime estimated costs.

We also conducted value of information analyses: a trial looking only at epidemiological parameters and costing up to £2.03 million could offer value for money. Our results indicated that most of the important uncertainty lies in the expected numbers of bleeds and strokes immediately following cAVR or Ross surgery; a trial estimating only these rates could be worth up to £1.90 million to decision makers. Limited value of £300 000–£425 000 was found for smaller study designs that measured utilities, poststroke disability cost or general event rates.

Our meta-analysis is in line with other recent findings, namely by Mazine *et al*³⁰ and McClure *et al*.³¹ Both found that the Ross procedure had generally better outcomes compared with biological AVR or mechAVR. McClure *et al*³¹ found lower thromboembolic events and better health-related quality of life for the Ross procedure, but neither meta-analysis considered cost-effectiveness generally or the value of further research.

As we rely on UK NHS costs, our cost-effectiveness and value of information results are specific to that perspective. However, the results of our meta-analysis and our modelling results of 15.2 (14.4 to 15.9) QALYs on the Ross procedure compared with 11.5 (10.4 to 13.3) QALYs on cAVR are potentially generalisable. To adapt to other countries, costs of events and management would need to be changed. However, when developing our model structure, we referred to models developed in the USA,

Table 3 EVPI and EVPPI for input parameters to the cost-effectiveness model

Parameters	Individual EVPPI	Population EVPPI*
Total EVPI	123.7 (118, 129.4)	2 026 008 (1 932 365; 2 119 651)
Meta-analysis	18.75 (3.887, 33.62)	307 068 (63 655; 550 481)
Utilities	25.97 (11.11, 40.83)	425 290 (181 877; 668 702)
Poststroke disability costs	20.94 (6.078, 35.81)	342 953 (99 541; 586 366)
Bleed or stroke following cAVR or Ross†	116.1 (101.2, 130.9)	1 900 959 (1 657 546; 2 144 372)
Other epidemiological parameters‡	31.15 (16.28, 46.01)	510 077 (266 665; 753 490)
Ross trial (no costs or utilities)§	123.7 (108.9, 138.6)	2 026 008 (1 782 596; 2 269 421)

*Population EVPPI given by multiplying individual EVPPI by 8263 population size.

†Early events of stroke or bleed following any of biological AVR, mechAVR or Ross procedure.

‡Other epidemiological parameters are probabilities of disability following stroke; stroke following reoperations for IE on AoV, reoperations for any cause on AoV, reoperations on the pulmonary valve; death following stroke, bleeding events, conservatively treated IE, reoperations for IE on AoV, reoperations for any cause on AoV, reinterventions on the pulmonary valve.

§Ross trial informing all parameters of the meta-analysis, bleed or stroke immediately following cAVR or Ross procedure, and other epidemiological parameters but not costs and utilities.

AoV, aortic valve; EVPI, expected value of perfect information; EVPPI, expected value of partial perfect information; IE, infective endocarditis; biological AVR, tissue aortic valve replacement; cAVR, conventional aortic valve replacement; mechAVR, mechanical aortic valve replacement.

Canada and Belgium,^{13 18 32} meaning it is not specific to the UK NHS. We would be willing to share our structure and software code with interested researchers to develop country-specific adaptations.

These results come at a time when the Ross operation is under much scrutiny, with several new studies reporting excellent long-term outcomes with this technique, achieving survival comparable with that of the general population.^{6 7} Several authors have suggested that the current guidelines regarding AVR in the young should be reconsidered with these new data in mind.^{33 34}

On the other hand, tissue valves are associated with poor results in the young, as shown in our meta-analysis, while mechanical valves bring the challenge of anticoagulation. Young patients, especially women planning a pregnancy, are reluctant to accept it, as it negatively impacts quality of life.³⁵ In addition, there is a cumulative risk of bleeding and thromboembolic events, shown to be significantly lower for Ross patients when compared with mechAVR.¹⁰ In our meta-analysis bleeding or thromboembolic events were four times more frequent after mechAVR, no doubt due to anticoagulation, with a non-negligible cumulative 1.55% rate per year. In younger, more active patients, the promise of fewer restrictions and lower risk of disability might overshadow the need for more reinterventions.

Given the more complex nature of the procedure, with more reinterventions being performed, there is the issue of the total burden on the health system, and whether the Ross procedure offers a sufficient advantage in quality of life to compensate for the added costs. Our economic analysis has shown that, although it has higher initial costs, the Ross procedure achieves a significant net benefit in the long term, no doubt through the significantly lower incidence of stroke.

At present, for the young patients with AoV disease, the European guidelines do not mention the Ross procedure

at all, while the American ones offer it as an alternative when anticoagulation is difficult to achieve.³⁶ The accumulated evidence in the past few years strongly suggests that the benefits from the Ross procedures outweigh the disadvantages in this group. Our study enforces this notion and adds the cost-effectiveness dimension. A clinical trial already exists comparing the Ross procedure with aortic homograft,⁸ but currently the main question is how it fares when tested against mechAVR and new-generation bioprostheses. A clinical trial to answer that question is warranted, with sufficient data coming from registries to support it; we have shown that even at a budget of over £1 million, it would still prove value for money to the health service providers.

Limitations

To design a computationally practical model, we did not include every adverse event associated with AVR. We excluded renal failure, pacemaker, arrhythmia and myocardial infarction, which were included in previous models,^{12–15} although these are less relevant in non-elderly patients. Our model recorded only history of the most serious event. These limitations could be overcome by expanding the number of model states, as in atrial fibrillation models,²⁴ or by including tunnel states,²⁶ but this could render EVPPI analysis computationally infeasible.³⁷ Adopting an NHS rather than a societal perspective limits our model's utility. As the Ross procedure is associated with a lower stroke and reoperation rate than biological AVR, including societal costs would likely increase its cost-effectiveness. Although there was limited uncertainty in our results, there were few high-quality studies available on many epidemiological parameters and utilities, which we represented with wide uncertainty margins. Further limitations arise from the meta-analysis used to inform the parameters and utilities of the

economic model. This was primarily based on observational studies, most of them retrospective. Selection criteria were used, and unpublished data, abstracts and presentations were excluded, generating a selection bias. There is also an inherent bias in the centres used for such studies, as the requirement for expertise in the Ross procedure limits the choice of centres. Finally, we have not estimated the expected value of sample information (EVSI). EVPI and EVPPI provide only an upper bound on trial value, while EVSI values specific trial designs.^{37 38} There is also substantial uncertainty about the number of Ross-eligible patients per year; our estimate of 960 patients per year in the UK was based on extrapolations from the existing evidence, although we were conservative in our estimates.

CONCLUSIONS

In young patients needing AVR, the Ross operation achieves better long-term survival, fewer bleeding and thromboembolic events, at the cost of more reinterventions, including those on the right ventricular outflow tract. Furthermore, our economic analysis showed that the Ross procedure is more cost-effective compared with cAVR. Despite this, the Ross operation does not have a clear place in guidelines, in lack of a good-quality clinical trial. Our study found that a trial comparing these options costing up to £2 million could offer value for money. An international collaboration started to generate this evidence prospectively and the trial is now in the pilot phase.

Acknowledgements We thank Umberto Benedetto for his initial input in the meta-analysis, Wei Fang for his advice on using multilevel Monte Carlo for EVPPI, and Ismail El-Hamamsy for advice on estimating the population eligible for Ross in the UK.

Contributors All authors contributed in a significant manner to the manuscript.

Funding SCS and JR received support from David Telling grant number 392 for this research. The Medical Research Council Hubs for Trials Methodology Research Collaboration and innovation in Difficult and Complex randomised controlled Trials In Invasive procedures (ConDuCT-II) provided support to HT, who was also supported by the NIHR Bristol Biomedical Research Centre. The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health. ACCV, EK, DMD, DF and MTAS confirm they have no financial support for this project to disclose for this research.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- Whitlock RP, McClure GR, Eikelboom JW. Aortic valve replacement in younger patients. *Eur Heart J* 2017;38:3378–81.
- Bourguignon T, Bouquiaux-Stablo A-L, Candolfi P, et al. Very long-term outcomes of the Carpentier-Edwards Perimount valve in aortic position. *The Annals of Thoracic Surgery* 2015;99:831–7.
- Stoica S, Goldsmith K, Demiris N, et al. Microsimulation and clinical outcomes analysis support a lower age threshold for use of biological valves. *Heart* 2010;96:1730–6.
- Kvidal P, Bergström PR, Hörte L-G, et al. Observed and relative survival after aortic valve replacement. *Journal of the American College of Cardiology* 2000;35:747–56.
- Forcillo J, El Hamamsy I, Stevens L-M, et al. The Perimount Valve in the Aortic Position: Twenty-Year Experience With Patients Under 60 Years Old. *The Annals of Thoracic Surgery* 2014;97:1526–32.
- Sharabiani MTA, Dorobantu DM, Mahani AS, et al. Aortic Valve Replacement and the Ross Operation in Children and Young Adults. *J Am Coll Cardiol* 2016;67:2858–70.
- Sievers H-H, Stierle U, Charitos EI, et al. A multicentre evaluation of the autograft procedure for young patients undergoing aortic valve replacement: update on the German Ross registry. *Eur J Cardiothorac Surg* 2016;49:212–8.
- El-Hamamsy I, Eryigit Z, Stevens L-M, et al. Long-term outcomes after autograft versus homograft aortic root replacement in adults with aortic valve disease: a randomised controlled trial. *The Lancet* 2010;376:524–31.
- Pibarot P, Dumesnil JG, Briand M, et al. Hemodynamic performance during maximum exercise in adult patients with the Ross operation and comparison with normal controls and patients with aortic bioprostheses. *Am J Cardiol* 2000;86:982–8.
- Mazine A, Ouzounian M. Aortic valve replacement in young and middle-aged adults: Looking beyond the tree that hides the forest. *Ann Transl Med* 2017;5.
- Welton NJ, Sutton AJ, Cooper NJ, et al. *Evidence synthesis for decision making in healthcare*. John Wiley and Sons, 2012.
- Brecker S, Mealing S, Padhiar A, et al. Cost-utility of transcatheter aortic valve implantation for inoperable patients with severe aortic stenosis treated by medical management: a UK cost-utility analysis based on patient-level data from the advance study. *Open Heart* 2014;1:e000155.
- Doble B, Blackhouse G, Goeree R, et al. Cost-effectiveness of the Edwards SAPIEN transcatheter heart valve compared with standard management and surgical aortic valve replacement in patients with severe symptomatic aortic stenosis: a Canadian perspective. *The Journal of Thoracic and Cardiovascular Surgery* 2013;146:52–60.
- Fairbairn TA, Meads DM, Hulme C, et al. The cost-effectiveness of transcatheter aortic valve implantation versus surgical aortic valve replacement in patients with severe aortic stenosis at high operative risk. *Heart* 2013;99:914–20.
- Orlando R, Pennant M, Rooney S, et al. Cost-effectiveness of transcatheter aortic valve implantation (TAVI) for aortic stenosis in patients who are high risk or contraindicated for surgery: a model-based economic evaluation. *Health Technol Assess* 2013;17:1–86.
- Murphy A, Fenwick E, Toff WD, et al. Transcatheter aortic valve implantation for severe aortic stenosis: the cost-effectiveness case for inoperable patients in the United Kingdom. *Int J Technol Assess Health Care* 2013;29:12–19.
- Siebert U, Alagoz O, Bayoumi AM, et al. State-Transition modeling: a report of the ISPOR-SMDM modeling good research practices task Force-3. *Value in Health* 2012;15:812–20.
- Gada H, Kapadia SR, Tuzcu EM, et al. Markov model for selection of aortic valve replacement versus transcatheter aortic valve implantation (without replacement) in high-risk patients. *The American Journal of Cardiology* 2012;109:1326–33.
- Fairbairn TA, Meads DM, Mather AN, et al. Serial change in health-related quality of life over 1 year after transcatheter aortic valve implantation. *Journal of the American College of Cardiology* 2012;59:1672–80.
- Reynolds MR, Lei Y, Wang K, et al. Cost-Effectiveness of Transcatheter Aortic Valve Replacement With a Self-Expanding Prosthesis Versus Surgical Aortic Valve Replacement. *Journal of the American College of Cardiology* 2016;67:29–38.
- Luengo-Fernandez R, Yin GSC, Gray AM, et al. Population-based study of acute- and long-term care costs after stroke in patients with AF. *International Journal of Stroke* 2013;8:308–14.
- Robinson A, Thomson R, Parkin D, et al. How patients with atrial fibrillation value different health outcomes: a standard gamble study. *J Health Serv Res Policy* 2001;6:92–8.
- Sterne JAC, Bodalia PN, Bryden PA, Caldwell D, et al. Oral anticoagulants for primary prevention, treatment and secondary prevention of venous thromboembolic disease, and for prevention of stroke in atrial fibrillation: systematic review, network meta-analysis and cost-effectiveness analysis. *Health Technol Assess* 2017;21:1–386.
- López-López JA, Sterne JAC, Thom HHZ, et al. Oral anticoagulants for prevention of stroke in atrial fibrillation: systematic review,

- network meta-analysis, and cost effectiveness analysis. *BMJ* 2017;82(8A).
25. Sterne JAC, Boudalia PN, Bryden PA, *et al.* Oral anticoagulants for primary prevention, treatment and secondary prevention of venous thromboembolic disease, and for prevention of stroke in atrial fibrillation: systematic review, network meta-analysis and cost-effectiveness analysis. *Health Technol Assess* 2017;21:1–386.
 26. Briggs AH, Claxton K, Sculpher MJ. *Decision modelling for health economic evaluation*. Oxford University Press: Oxford, 2006.
 27. Neumann PJ, Cohen JT, Weinstein MC. Updating Cost-Effectiveness — The Curious Resilience of the \$50,000-per-QALY Threshold. *N Engl J Med* 2014;371:796–7.
 28. Giles MB, Goda T. Decision-making under uncertainty: using MLMC for efficient estimation of EVPPI. *ArXiv* 2017;05531.
 29. Weber A, Nouredine H, Englberger L, *et al.* Ten-year comparison of pericardial tissue valves versus mechanical prostheses for aortic valve replacement in patients younger than 60 years of age. *The Journal of Thoracic and Cardiovascular Surgery* 2012;144:1075–83.
 30. Mazine A, Rocha RV, El-Hamamsy I, *et al.* Ross procedure vs mechanical aortic valve replacement in adults: a systematic review and meta-analysis. *JAMA Cardiol* 2018;3.
 31. McClure GR, Belley-Cote EP, Um K, *et al.* The Ross procedure versus prosthetic and homograft aortic valve replacement: a systematic review and meta-analysis. *Eur J Cardiothorac Surg* 2019;55:247–55.
 32. Neyt M, Van Brabant H, Devriese S, *et al.* A cost-utility analysis of transcatheter aortic valve implantation in Belgium: focusing on a well-defined and identifiable population. *BMJ Open* 2012;2.
 33. El-Hamamsy I, Bouhout I. The Ross procedure: time for a hard look at current practices and a reexamination of the guidelines. *Ann. Transl. Med.* 2017;5.
 34. David TE. Aortic valve replacement in children and young adults. *J Am Coll Cardiol* 2016;67:2871–3.
 35. Nötzold A, Hüppe M, Schmidtke C, *et al.* Quality of life in aortic valve replacement: pulmonary autografts versus mechanical prostheses. *Journal of the American College of Cardiology* 2001;37:1963–6.
 36. Nishimura RA, Otto CM, Bonow RO, *et al.* 2017 AHA/ACC focused update of the 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. *Circulation* 2017;135:e1159–95.
 37. Welton NJ, Thom HHZ. Value of information: we've got speed, what more do we need? *Med Decis Making* 2015;35:564–6.
 38. Welton NJ, Madan JJ, Caldwell DM, *et al.* Expected value of sample information for multi-arm cluster randomized trials with binary outcomes. *Med Decis Making* 2014;34:352–65.

Correction: *Clinical and cost-effectiveness of the Ross procedure versus conventional aortic valve replacement in young adults*

Thom H, Visan AC, Keeney E, *et al.* Clinical and cost-effectiveness of the Ross procedure versus conventional aortic valve replacement in young adults. *Open Heart* 2019;6:e001047. doi:10.0036/openhrt-2019-001047

The authors want to alert readers to the following error identified in the published version.

The co-author name Alexandru Ciprian Ciprian Visan has been corrected to Alexandru Ciprian Visan

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

Open Heart 2019;6:e001047corr1. doi:10.1136/openhrt-2019-001047corr1