


openheart ASPIRE-3-PREVENT: a cross-sectional survey of preventive care after a coronary event across the UK

Catriona S Jennings ^{1,2}, Kornelia Kotseva,^{1,2} Paul Bassett,¹ Agnieszka Adamska,^{1,2} David Wood,^{1,2} on behalf of the ASPIRE-3-PREVENT Investigators

To cite: Jennings CS, Kotseva K, Bassett P, *et al.* ASPIRE-3-PREVENT: a cross-sectional survey of preventive care after a coronary event across the UK. *Open Heart* 2020;**7**:e001196. doi:10.1136/openhrt-2019-001196

Received 24 October 2019
Revised 3 February 2020
Accepted 26 February 2020

ABSTRACT

Objective To quantify the implementation of the third Joint British Societies' Consensus Recommendations for the Prevention of Cardiovascular Disease (JBS3) after coronary event.

Methods Using a cross-sectional survey design, patients were consecutively identified in 36 specialist and district general hospitals between 6 months and 2 years, after acute coronary syndrome or revascularisation procedure and invited to a research interview. Outcomes included JBS3 lifestyle, risk factor and therapeutic management goals. Data were collected using standardised methods and instruments by trained study nurses. Blood was analysed in a central laboratory and a glucose tolerance test was performed.

Results 3926 eligible patients were invited to participate and 1177 (23.3% women) were interviewed (30% response). 12.5% were from black and minority ethnic groups. 45% were persistent smokers, 36% obese, 52.9% centrally obese, 52% inactive; 30% had a blood pressure >140/90 mm Hg, 54% non-high-density lipoprotein ≥ 2.5 mmol/L and 44.3% had new dysglycaemia. Prescribing was highest for antiplatelets (94%) and statins (85%). 81% were advised to attend cardiac rehabilitation (86% <60 years vs 79% ≥ 60 years; 82% men vs 77% women; 93% coronary artery bypass grafting vs 59% unstable angina), 85% attended if advised; 69% attended overall. Attenders were significantly younger ($p=0.03$) and women were less likely to attend ($p=0.03$).

Conclusions Patients with coronary heart disease (CHD) are not being adequately managed after event with preventive measures. They require a structured preventive cardiology programme addressing lifestyle, risk factor management and adherence to cardioprotective medications to achieve the standards set by the British Association for Cardiovascular Prevention and Rehabilitation and JBS3 guidelines.

INTRODUCTION

ASPIRE-3-PREVENT (A-3-P)^{1,2} was conducted to evaluate preventive care for patients with coronary heart disease (CHD) in everyday clinical practice. Goals for care were outlined, most recently, in the Joint British Societies' Consensus Recommendations for the

Key questions

What is already known about this subject?

► There is strong scientific evidence that lifestyle, risk factor and therapeutic management of patients with CHD will reduce their risk of subsequent events and increase survival. The Joint British Societies' Consensus Recommendations for the Prevention of Cardiovascular Disease guidelines summarise this scientific evidence for secondary prevention and the British Association for Cardiovascular Prevention and Rehabilitation advocate standards for cardiovascular disease prevention and rehabilitation.

What does this study add?

► ASPIRE-3-PREVENT provides a contemporary picture of the implementation of the guidelines and standards in everyday clinical practice across the UK and quantifies the scope for improvement.

How might this impact on clinical practice?

► These results will inform commissioners of services about the need to improve delivery of secondary preventive care and how this can be achieved through a modern preventive cardiology programme.

Prevention of Cardiovascular Disease (JBS3)³ and aim to reduce morbidity, improve quality of life and increase life expectancy. A-3-P was conducted as part of the fifth EUROASPIRE survey, which investigated preventive care in patients with CHD from 27 European countries. The European Society of Cardiology (ESC) EUROASPIRE cross-sectional surveys⁴⁻⁸ have repeatedly evaluated implementation of the European prevention guidelines since 1995.

There is substantial evidence that intervening to modify adverse lifestyle behaviours and manage medical risk factors can reduce the risk of future cardiovascular events and improve survival in patients with CHD.^{3,9-11} Opportunities for delivering quality care for this patient group are offered by comprehensive cardiovascular prevention and



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹NHLI, Imperial College London, London, UK

²National Institute for Prevention and Cardiovascular Health, National University of Ireland Galway, Galway, Ireland

Correspondence to

Dr Catriona S Jennings; catriona.jennings@nuigalway.ie

rehabilitation programmes. In the third edition of their standards and core components,¹² the British Association for Cardiovascular Prevention and Rehabilitation (BACPR) place emphasis on the need to underpin cardiovascular disease (CVD) risk reduction with patient-centred, behavioural and goal-oriented approaches to address risk factor and therapeutic management with timely intervention from a clinically led multidisciplinary team. Audit of outcomes is through the National Audit for Cardiac Rehabilitation's (NACR)¹³ latest report 2018.

The specific objectives of this survey were:

- ▶ To determine adherence to JBS3 guidelines in everyday clinical practice for patients with CHD.
- ▶ To evaluate diagnostic and therapeutic strategies for dysglycaemia.
- ▶ To make recommendations to the BACPR, British Cardiovascular Society and Public Health England on ways to improve preventive cardiology care.

This paper describes the principal results of A-3-P in relation to achievement of lifestyle, risk factor and therapeutic targets as defined in the JBS3 guidelines.

METHODS

A-3-P was registered as a cross-sectional survey with the National Institute for Health Research Clinical Research Network (CRN) Portfolio on 23 August 2016 and made visible to all CRNs in England wishing to participate. It aimed to secure representation from hospital centres across five English National Health Service regions and in CRNs in Wales, Northern Ireland and Scotland. Thirty-six specialist cardiac centres and district general hospitals were recruited in this convenience sample.

Fieldwork for the survey was led by CRN and hospital Research and Development (R&D) department nurses. Hospitals were asked to create lists of retrospectively identified eligible patients from their clinical coding systems and to recruit consecutively according to admission date for the recruiting event. Eligibility was defined as male and female patients between the ages of 18 and 79 years at the time of diagnosis with new or recurrent ST-elevation myocardial infarction (STEMI), non-STEMI (NSTEMI) or acute ischaemia and/or elective or emergency percutaneous coronary intervention (PCI) or coronary artery bypass grafting (CABG). Patients were entered onto an electronic log and invited to participate, where reason for non-participation was also recorded. Patients were invited to attend an interview and examination with the centrally trained nurses who obtained written, informed consent, collected data using standardised instruments and methods (see [box 1](#)), and drew blood, processed and stored it prior to it being sent to the central laboratory in Finland for analysis. Outcome measures can be seen in [box 1](#). Data were entered and managed on the anonymised ESC EurObservational Research Programme electronic online database.

A sample size of 400 patients attending for interview was sufficient with a precision of at least 5% and a confidence

Box 1 Outcome measures and research instruments for the ASPIRE-3-PREVENT (A-3-P) survey

- ▶ Current smoking validated with a breath carbon monoxide (CO) test ≥ 10 ppm (Micro+Smokerlyzer, Bedfont Scientific, Kent, UK).
- ▶ Persistent smoking: smoking at recruiting event and still smoking at interview.
- ▶ Overweight^{26 27}: body mass index (BMI) ≥ 25 kg/m² (non-Asian); BMI ≥ 23 kg/m² (Asian) (height and weight measured in light indoor clothes without shoes with SECA measuring stick model 220 and SECA scale model 701).
- ▶ Obesity^{26 27}: BMI ≥ 30 kg/m² (non-Asian); BMI ≥ 25 kg/m² (Asian).
- ▶ Central overweight^{26 27}: waist circumference (WC) ≥ 94 cm (non-Asian men); WC ≥ 90 cm (Asian men); WC ≥ 80 cm (all women) (WC measured with metal tape measures mid-way between the lower rib margin and the iliac crest).
- ▶ Central obesity^{26 27}: WC ≥ 102 cm (non-Asian men); WC ≥ 90 cm (Asian men); WC ≥ 88 cm (non-Asian women); WC ≥ 80 cm (Asian women).
- ▶ Not performing moderate-intensity physical activity at least 30 min/5 times per week (Godin Leisure-Time Exercise Questionnaire²⁸).
- ▶ Elevated blood pressure: ≥ 140 mm Hg and/or ≥ 90 mm Hg (blood pressure measured twice on the right upper arm in a sitting position after rest for at least 5 min using Omron M6 HEM 7211-E automatic digital sphygmomanometers).
- ▶ Non-high-density lipoprotein (HDL) cholesterol: ≥ 2.5 mmol/L (analysed in central laboratory in Finland).
- ▶ Low-density lipoprotein cholesterol (LDL-C) ≥ 1.8 mmol/L.
- ▶ HbA1c $\geq 7\%$ (≥ 53 mmol/mol) in patients with type 2 diabetes mellitus.²⁹
- ▶ New diabetes mellitus (DM); impaired fasting glycaemia (IFG); impaired glucose tolerance (IGT) from oral glucose tolerance test (OGTT) (see [box 2](#)³⁰) in patients without self-reported diabetes and with fasting blood glucose (FBG) < 11.1 mmol/L measured with HemoCue near-patient testing monitor.
- ▶ Cardiovascular drug therapies and medication adherence.³¹
- ▶ Hospital Anxiety and Depression Scale (HADS³²), HeartQoL,³³ EuroQoL (EQ-5D³⁴).

of 95% to estimate prevalence and allow power to stratify by age and gender and to estimate prevalence of single risk factors with 95% CIs of width 10%. All statistical analyses were undertaken using STATA V.11 (StataCorp). The majority of analyses were descriptive aiming to summarise characteristics and outcomes. The exact binomial method was used to calculate CIs for the percentage values. The McNemar test was used to statistically compare the results from hospital discharge and the time of interview for the proportion of patients on the most popular medications.

The cardiac rehabilitation (CR) analyses focused on differences in patients who were advised and not advised and who attended and did not attend a programme. Attendance was defined as one or more of the programme sessions. The groups were compared on key demographic variables and comparisons between the groups were made for lifestyle, risk factors and prescribing. All outcomes were binary and comparisons between groups were made using logistic regression. Adjustments were made if results varied in terms of key demographics.

Box 2 Diagnostic criteria for diabetes and dysglycaemia**Diabetes** No Yes

Fasting plasma glucose ≥ 7.0 mmol/L (126 mg/dL)
or

2-hour plasma glucose* ≥ 11.1 mmol/L (200 mg/dL)

Impaired glucose tolerance (IGT) No Yes

Fasting plasma glucose < 7.0 mmol/L (< 126 mg/dL)
and 2-hour glucose* ≥ 7.8 and < 11.1 mmol/L (140 and < 200 mg/dL)

Impaired fasting glucose (IFG) No Yes

Fasting plasma glucose ≥ 6.1 and < 6.9 mmol/L (110 and < 125 mg/dL)

and 2-hour plasma glucose* < 7.8 mmol/L (140 mg/dL)

*Venous plasma glucose 2 hours after the ingestion of 75 g dissolved glucose in 200 mL water.

RESULTS

Figure 1 shows the flow of patients through the study. Non-attendance at interview was driven principally by a failure to respond to the invitation (71% of DNA). As expected in a population of patients with CHD under the age of 80 years, the majority were male (77%). The mean age of the participants was 65.7 years and most were of white ethnicity. Nearly two-thirds of patients identified for the study had either acute STEMI or NSTEMI.

Table 1 shows lifestyle habits, medical risk factors, prescribing and psychosocial status in all patients and also according to their attendance at a CR programme. If adhering to national standards (BACPR), CR should have been offered to all patients who have suffered a cardiac event and/or who have been revascularised, that is, all those who were recruited to this survey. In fact, a large proportion of interviewed patients (80.8%) were advised to attend a programme and, of these, 84.8% attended. Out of all 1177 patients, 68.5% attended a programme. Patients most likely to be advised were those who had undergone CABG. Least advised were patients with unstable angina, older patients, women, those not achieving the physical activity target and those with self-reported diabetes. Interestingly, less advised groups were just as likely to attend if so advised (see table 2). There were significant differences between patients who attended and did not attend in relation to mean age (66.6 years in non-attenders compared with 65.4 years in attenders, $p=0.03$), gender and revascularisation procedure (see table 2) with women and older patients being less likely to attend and those revascularised with CABG and acute myocardial infarction being more likely to attend.

Tobacco

Around one-fifth reported smoking at the time of their event (table 1) and nearly one-half of these were still smoking at the time of interview. There were no statistical differences between men and women with regard to smoking habit. Among current smokers, two-thirds

reported an intention to quit within the next 6 months. There were no gender differences.

While the majority of current smokers or recent quitters reported being advised to quit since the recruiting cardiac event, only a minority were advised to use pharmacological support or referred for specialist help (see table 3). At the time of discharge from hospital, around one-fifth of current smokers and recent quitters were prescribed antismoking treatments (table 4). Around two-thirds of patients made a quit attempt although a relatively small proportion accessed specialist support or pharmacotherapies. Current smokers were significantly less likely to attend a CR programme.

Overweight and obesity

A large majority of patients were above ideal body mass index (table 1). More than half were centrally obese with a significantly higher prevalence in women. There




Figure 1 Flow chart of patients through all phases of data collection in ASPIRE-3-PREVENT. CABG, coronary artery bypass grafting; NSTEMI, non-STEMI; OGTT, oral glucose tolerance test; PTCA, percutaneous transluminal coronary angioplasty; STEMI, ST-elevation myocardial infarction.

Table 1 Lifestyle and risk factor outcomes in patients following an acute cardiac event comparing those who attended and those who did not attend cardiac rehabilitation

	Attended cardiac rehabilitation n=800				Did not attend cardiac rehabilitation n=368				All patients** n=1177			
	M	F	All	n (%)	M	F	All	n (%)	M	F	All	n (%)
	629	171	800		268	100	368		903	274	1177	
Current smoker	48 (7.6)	7 (4.1)	55 (6.9)	32 (11.9)	11 (11.0)	43 (11.7)*	83 (9.2)	19 (6.9)	102 (8.7)			
Smoker at index event	107 (17.0)	32 (18.7)	139 (17.4)	53 (19.8)	20 (20.0)	73 (19.8)	163 (18.1)	53 (19.3)	216 (18.4)			
Persistent smoker†	46 (43.0)	7 (21.9)	53 (38.1)	30 (56.6)	11 (55.0)	41 (56.2)	79 (48.5)	19 (35.9)	98 (45.4)			
Not achieving physical activity guidelines	278 (45.1)	103 (62.1)	381 (48.7)	149 (58.2)	64 (64.7)	213 (60.0)**	428 (48.9)	169 (63.1)***	597 (52.2)			
Overweight	488 (77.7)	126 (74.1)	614 (76.9)	213 (79.8)	83 (83.0)	296 (80.7)	705 (78.4)	212 (77.7)	917 (78.2)			
Obese	221 (35.2)	62 (36.5)	283 (35.5)	95 (35.6)	43 (43.0)	138 (37.6)	316 (35.2)	106 (38.8)	422 (36.0)			
Centrally overweight	446 (71.4)	146 (85.9)	592 (74.5)	192 (72.5)	85 (87.6)	277 (76.5)	671 (71.7)	234 (86.7)***	875 (75.2)			
Centrally obese	296 (47.4)	105 (61.8)	401 (50.4)	135 (50.9)	76 (78.4)	211 (58.3)	432 (48.3)	184 (68.2)***	616 (52.9)			
BP ≥140/90 mm Hg	181 (28.8)	46 (26.9)	227 (28.4)	92 (34.3)	30 (30.0)	122 (33.2)	275 (30.5)	79 (28.8)	354 (30.1)			
BP ≥130/80 mm Hg	363 (57.7)	89 (52.1)	452 (56.5)	158 (59.0)	58 (58.0)	216 (58.7)	525 (58.2)	150 (54.7)	675 (57.4)			
LDL-C ≥1.8 mmol/L	334 (56.4)	110 (66.7)	444 (58.7)	163 (69.9)	66 (74.2)	229 (66.6)†	500 (58.7)	177 (68.9)**	677 (61.0)			
Non-HDL-C ≥2.5 mmol/L	292 (48.6)	103 (62.1)	395 (51.5)	143 (55.2)	65 (69.2)	208 (58.9)†	439 (50.7)	169 (64.3)	609 (53.9)			
Self-reported diabetes	127 (20.3)	34 (20.1)	161 (20.3)	79 (29.5)	23 (23.0)	102 (27.7)	209 (23.3)	58 (21.3)	267 (22.8)			
HbA1c ≥7% (>53 mmol/mol in self-reported diabetes)	47 (7.9)	18 (10.8)	65 (8.5)	43 (16.7)	11 (11.7)	54 (15.4)	92 (10.7)	29 (11.0)	121 (8.8)			
OGTT: IGT	122 (19.7)	39 (23.4)	161 (20.5)	31 (11.7)	20 (20.2)	51 (14.0)	153 (17.2)	59 (21.9)	212 (18.3)			
IFG	79 (12.8)	18 (10.8)	97 (12.3)	25 (9.4)	8 (8.1)	33 (9.0)	104 (11.7)	27 (10.0)	131 (11.3)			
New diabetes	94 (15.2)	21 (12.6)	115 (14.6)	44 (16.5)	11 (11.1)	55 (15.1)	138 (15.5)	32 (11.9)	170 (14.7)			
Antiplatelet	603 (95.9)	165 (96.5)	768 (96.0)	243 (91.0)	88 (88.0)	331 (90.2)*	849 (94.4)	256 (93.4)	1105 (94.2)			
Beta blocker	503 (80.1)	129 (75.4)	632 (79.1)	199 (75.1)	68 (68.0)	267 (73.2)	704 (78.6)	198 (72.3)	902 (77.1)			
ACE	402 (63.9)	87 (50.9)	489 (61.1)	160 (59.9)	53 (53.0)	213 (58.0)	565 (62.9)	142 (51.8)	707 (60.3)			
ARB	98 (15.6)	41 (24.0)	139 (17.4)	36 (13.5)	18 (18.0)	54 (14.8)	134 (14.9)	60 (21.9)	194 (16.6)			
Other BP-lowering drugs	166 (26.4)	51 (29.8)	217 (27.1)	93 (34.8)	34 (34.0)	127 (34.6)	261 (29.0)	86 (31.4)	347 (29.6)			
Statins	545 (86.8)	143 (83.0)	687 (86.0)	225 (84.6)	75 (75.0)	300 (82.0)	773 (86.2)	220 (80.3)	993 (84.8)			
Other lipid-lowering drugs	29 (4.6)	9 (5.3)	38 (4.8)	16 (6.0)	5 (5.0)	21 (5.7)	45 (5.0)	14 (5.1)	59 (5.0)			
Drug adherence§												
BP lowering	584/588 (99.3)	158/159 (99.4)	742/747 (99.3)	95/101 (94.1)	30/31 (96.8)	125/132 (94.7)	818/835 (98.0)	244/246 (99.2)	1062/1081 (98.2)			
Lipid lowering	528/532 (99.3)	149/150 (99.3)	677/682 (99.3)	91/93 (97.9)	28/29 (96.6)	119/122 (97.5)	744/756 (98.4)	232/234 (99.2)	990/976 (98.6)			

Continued

Table 1 Continued

	Attended cardiac rehabilitation n=800			Did not attend cardiac rehabilitation n=368			All patients** n=1177		
	M	F	All	M	F	All	M	F	All
	629	171	800	268	100	368	903	274	1177
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Glucose lowering	169/178 (94.9)	46/47 (97.9)	215/225 (95.6)	34/35 (97.1)	10/11 (90.9)	44/46 (95.7)	261/273 (95.6)	74/76 (97.4)	335/349 (96.0)
HADS Anxiety ≥8	123 (19.8)	54 (32.3)	177 (22.4)	51 (19.5)	39 (39.8)	90 (25.1)	176 (19.8)	94 (35.1)	270 (23.4)
HADS Depression ≥8	75 (12.1)	17 (10.2)	92 (11.7)	38 (14.6)	24 (24.4)	62 (17.3)	113 (12.7)	41 (15.3)	154 (13.3)
EQ-VAS out of 100†	80 (70, 90)	80 (60, 90)	80 (70, 90)	80 (65, 90)	75 (50, 90)	76 (60, 90)	80 (70, 90)	75 (60, 90)	80 (68, 90)

*P=0.006; **p=0.003; ***p<0.001.

†P=0.02.

‡Persistent smoker: smoking at hospital admission and still smoking at interview.

§Medication adherence denominator included due to variation in drug treatment.

¶Median (IQR).

**No data available on CR attendance in 9 patients

ARB, angiotensin receptor blocker; BP, blood pressure; EQ-VAS, EuroQol Visual Analogue Scale; HADS, Hospital Anxiety and Depression Scale; HDL-C, high-density lipoprotein cholesterol; IFG, impaired fasting glycaemia; IGT, impaired glucose tolerance; LDL-C, low-density lipoprotein cholesterol; OGTT, oral glucose tolerance test.

were no significant differences in prevalence according to attendance at CR. Two-thirds of overweight patients reported being advised to lose weight and attempted to do so.

Physical activity and exercise

Approximately one-half reported not achieving the physical activity recommendation (see table 1). However, 40%, mostly women, reported having a disability of which the majority said limited exercise capacity (see table 3). Around one-half claimed to take regular planned exercise (see figure 2).

Biological risk factors

One-third had raised blood pressure (BP) according to the JBS3 goal (<140 and/or 90 mm Hg) (table 1). However, if more recent guidelines on BP targets from the American Heart Association/American College of Cardiology¹⁴ and the ESC/European Society of Hypertension¹⁵ are taken into account, this proportion is higher with nearly 60% of patients having a BP of ≥130 mm Hg and/or ≥80 mm Hg. More than one-half of patients are not reaching the JBS3 non-high-density lipoprotein cholesterol (HDL-C) goal of 2.5 mmol/L and these prevalences are significantly higher in women. While one-quarter of patients reported having been previously diagnosed with diabetes, an oral glucose tolerance test (OGTT) picked up new dysglycaemia in a further 44% of patients (see figure 3). 39.7% of patients with self-reported diabetes had a BP <130 mm Hg and/or <80 mm Hg and 50.9% an HbA1c of <7% (<53 mmol/mol). There were no gender differences in relation to BP control; however, low-density lipoprotein cholesterol (LDL-C) and non HDL-C were better managed in men (see table 5).

Prescribing

Prescribing (table 1) was, on the whole, very good with 85% or more of patients reporting taking antiplatelets, statins and BP-lowering drugs. However, despite more than 90% of patients prescribed BP-lowering drugs, less than three-quarters were at the JBS3 goal. This proportion was much lower (43.2%) if a target of 130/80 mm Hg is used, which is in keeping with recent guidelines.^{14 15} There is a similar picture for lipid control with less than half at goal despite 83% of patients being prescribed lipid-lowering drugs. In patients with self-reported diabetes at interview, glycaemic control was not at target in one-third of patients. Statins were less frequently prescribed in women (86.2% men, 80.3% women, p=0.02).

DISCUSSION

Our A-3-P survey investigated the implementation of CVD prevention guidelines in the UK and continues to highlight the enormous potential to reduce the risk of recurrent disease and associated disabilities and also to improve survival. In this discussion, we focus on achievement of a healthy lifestyle, treating risk factors—BP, lipids and

diabetes—to targets with cardioprotective medications, and the role of prevention and rehabilitation programmes.

Persistent smoking in those who were smoking at the time of their event is high at 45%. There is a lack of support in hospital and after discharge to help patients quit smoking or to maintain abstinence due to underprescribing of drugs and provision of counselling to support smoking cessation. Many of these persistent smokers intend to quit so there is the opportunity for a behavioural intervention supported by pharmacotherapies with proven efficacy,^{16 17} effectiveness¹⁸ and safety¹⁹ in this patient population. Some patients were prescribed nicotine replacement therapy at the time of discharge from hospital but what is required is a systematic evidence-based approach which starts on the first day of hospital admission and continues after discharge in the CR programme and in general practice. The recent announcement by the Department of Health to support smokers admitted to hospital to quit will improve smoking cessation rates including for those with coronary disease.²⁰

Encouragingly, more than half of the interviewed patients reported taking regular exercise and one-fifth report that they intended to start doing so within the next 6 months. Taking regular exercise was more common in men; however, the older age of female patients and their associated disabilities should be taken into account. Interestingly, patients not achieving the physical activity target were less likely to be advised to attend CR, and were also less likely to attend a programme, and yet these patients would derive even greater benefits from supervised exercise sessions.

There is a high prevalence of obesity and central obesity in this population, particularly in women. About two-thirds were advised to lose weight, which means that such advice was not given to a significant minority, and a similar proportion reported they were attempting to do so. One caveat is that for those who successfully quit smoking, weight gain is common²¹ and this may be a confounding factor in this overall picture.

The high prevalence of central obesity is a major factor in the increasing prevalence of diabetes in patients with CHD. Diabetes is commonly associated with coronary disease and, as the results of the OGTT demonstrate there is a considerable overlap with two-thirds of patients with CHD exhibiting some form of dysglycaemia. What is of great concern is that two-thirds of these patients with dysglycaemia had not been identified at the time of their event or subsequently. Diabetes is a major risk factor for coronary disease and patients with both CHD and diabetes are at much greater risk of a further major adverse cardiovascular event. So why are we not screening the population with CHD without self-reported diabetes with an OGTT, the only test which most reliably identifies all patients with diabetes and with impaired glucose tolerance?²² The identification of dysglycaemia will be a stimulus to greater diligence in monitoring for the development of diabetes, and in those with newly diagnosed diabetes, appropriate monitoring for the development of microvascular disease and other complications can be put in place.

Although cardioprotective medications are prescribed in all patients, the extent to which risk factors are controlled to national targets remains disappointing. Just over two-thirds of patients are achieving the conservative BP target of <140 mm Hg and/or 90 mm Hg, but only 42% are achieving the target of <130 mm Hg and/or 80 mm Hg. The non-HDL-C target of <2.5 mmol/L is achieved by just over half of all patients and only 44% are below the more stringent LDL-C target of <1.8 mmol/L. In patients with self-reported diabetes an HbA1c <7.0% is achieved in just over half of all patients. In these patients with diabetes the lower BP target of <130 mm Hg and/or 80 mm Hg is only achieved by just over a third.

After experiencing an acute cardiac event and/or undergoing a revascularisation procedure, patients require an interdisciplinary CR programme which provides expert

Table 2 Characteristics of patients advised to attend, participated if advised and who participated overall in a cardiac rehabilitation programme

Patient group	Advised to participate		Participated if advised		Participated overall	
	n/N (%)	P value	n/N (%)	P value	n/N (%)	P value
All patients	944/1168 (80.8)		800/944 (84.8)		800/1168 (68.5)	
Age <60	254/296 (85.8)	0.01	213/254 (83.9)	0.65	213/296 (72.0)	0.14
Age ≥60	690/872 (79.1)		587/690 (85.1)		587/872 (67.3)	
Male	735/897 (81.9)	0.08	629/735 (85.6)	0.18	629/897 (70.1)	0.03
Female	209/271 (77.1)		171/209 (81.8)		171/271 (63.1)	
AMI (STEMI)	213/242 (88.0)	<0.001	180/213 (84.5)	0.15	180/242 (74.4)	<0.001
AMI (non-STEMI)	428/514 (83.3)		356/428 (83.2)		356/514 (69.3)	
CABG	99/107 (92.5)		87/99 (87.9)		87/107 (81.3)	
PTCA	153/218 (70.2)		128/153 (83.7)		128/218 (58.7)	
Unstable angina	51/87 (58.6)		49/51 (96.1)		49/87 (56.3)	

AMI, acute myocardial infarction; CABG, coronary artery bypass grafting; PTCA, percutaneous transluminal coronary angioplasty; STEMI, ST-elevation myocardial infarction.

Table 3 Lifestyle habits and change attempts in patients participating in the A-3-P survey

	Men %	Women %	All %
Tobacco			
Referred to specialist clinic	32.7	40.4	34.6
Advised to stop			
Use NRT	34.4	50.0 (p=0.04)	38.1
Use bupropion	2.6	2.0	2.5
Use varenicline	5.9	3.9	5.4
Made quit attempt	60.1	71.7	63
Tried to reduce	37.5	26.4	34.7
Attended specialist clinic	14.4	11.3	13.6
Used NRT	23.8	24.5	23.9
Used bupropion	0.0	1.9	0.5
Used varenicline	3.3	1.9	2.9
Diet and weight			
Dietary advice	75.4	71.5	74.5
Advised to lose weight	63.7	59.4	62.7
Steps taken—improve diet			
Reduce salt	67.2	60.5 (p=0.04)	65.7
Reduce fat	69.2	69.5	69.2
Change fat	61.6	60.3	61.3
Increase fruits and vegetables	72.9	65.7 (p=0.02)	71.2
Eat more fish	61.3	56.4	60.1
Reduce sugar	64.7	55.9 (p=0.009)	62.6
Reduce alcohol	51.6	40.7 (p=0.002)	49.1
Eat more sterols	29.4	17.1 (p<0.001)	26.5
Steps taken to lose weight—in overweight patients	67.6	65.6	67.1
Physical activity			
Disability/infirmity	38.1	51.1 (p<0.001)	41.1
Infirmity limits activity	74.6	85 (p=0.01)	77.6
Professional advice to increase activity	52.4	57.1	53.5
Advised to increase everyday activities	60.6	59.9	17.2
Advised to join walking group	16.8	18.3	68.8
Any steps taken to increase activity	70.7	71.2	70.8

A-3-P, ASPIRE-3-PREVENT; NRT, nicotine replacement therapy.

support to help patients and families change adverse health behaviours, to monitor and manage medical risk factors to targets and promote adherence to prescribed medications over the long term. Such a programme should be an integral part of their care with patients moving seamlessly from acute care to longer term preventive care to further reduce their risk of events, the need for further revascularisation and to improve survival.

Table 4 Prescription of tobacco cessation drugs at discharge and at interview

Time point	Treatment	Males n (%)	Females n (%)	All n (%)
Discharge (from medical notes)	NRT	27 (17.8)	14 (28.6)	41 (20.4)
	Bupropion	1 (0.7)	1 (2)	2 (1)
	Varenicline	1 (0.7)	1 (2)	2 (1)
Interview	NRT	10 (6.1)	1 (1.9)	11 (5.1)
	Bupropion	0 (0)	0 (0)	0 (0)
	Varenicline	0 (0)	0 (0)	0 (0)

NRT, nicotine replacement therapy.

In this survey, 68% participated in some form of CR programme by comparison with 47% reported by NACR, and therefore our patient sample is skewed towards those undertaking rehabilitation. Participation was higher for younger patients, men and those who had a CABG. Although these patients all had coronary atherosclerosis there was a large differential in participation rates from 81% for CABG to 56% for those presenting with acute myocardial ischaemia. Older patients and women were less likely to be advised to attend a programme and the same was true for patients with acute myocardial ischaemia. There should be no difference in participation rates by diagnostic category or gender given that they all have the same underlying disease.

A recent systematic review and meta-analysis of trials since 2010 of prevention and rehabilitation programmes showed no overall benefit in terms of all-cause mortality, but significant reductions in cardiovascular mortality, including stroke.²³ However, comprehensive programmes managing six or more risk factors did reduce all-cause mortality, compared with programmes managing less




Figure 2 Self-reported regular planned exercise.




Figure 3 Glucose metabolism in 1177 patients based on an oral glucose tolerance test in those without self-reported diabetes.

than six risk factors. Programmes that took responsibility for prescribing, uptitrating and monitoring adherence to cardioprotective medications also reduced all-cause

mortality compared with programmes that devolved that responsibility to other healthcare professionals. These results support the view that prevention and rehabilitation programmes should be truly comprehensive, addressing lifestyle, risk factor management and taking responsibility within the service provided to manage all cardioprotective medications.^{24 25}

Exercise-based CR is not truly comprehensive based on the results of this survey. Achievement of physical activity guidelines is better in those attending CR but there is no impact on persistent smoking, the prevalence of obesity and central obesity or control of BP to defined targets. Achievement of the LDL-C goal is significantly better in those attending CR but even so more than half of these patients are not achieving this target. Diabetes control is no better in those attending CR and new diabetes was missed just as frequently in those attending CR compared with those who did not. Cardioprotective medications were prescribed as commonly in those attending and not attending CR apart from antiplatelet medications which were significantly higher in those attending CR. Anxiety and depression outcomes were no different and health-related quality of life was the same. To reduce the risk of a further event and improve life expectancy, it is essential to address all aspects of lifestyle, manage the biological risk factors to targets and ensure adherence with optimally prescribed cardioprotective medications.

Table 5 Therapeutic control of blood pressure, cholesterol and glucose by gender

	Male (n=903)		Female (n=274)		P value
	n	% (95% CI)	n	% (95% CI)	
Blood pressure*	899		258		
BP at target 1†	595	69.9 (66.7 to 73.0)	187	72.4 (66.6 to 77.8)	0.43
BP at target 2‡	360	42.3 (39.0 to 45.7)	119	46.1 (39.9 to 52.4)	0.28
Cholesterol§	759		219		
TC <5.0 mmol/L	698	92.0 (89.8 to 93.8)	184	84.0 (78.5 to 88.6)	<0.001
TC <4.0 mmol/L	523	68.9 (65.4 to 72.2)	104	47.5 (40.7 to 54.3)	<0.001
LDL <1.8 mmol/L¶	329	43.9 (40.3 to 47.6)	75	34.7 (28.4 to 41.5)	0.02
LDL <2.0 mmol/L¶	441	58.9 (55.3 to 62.4)	93	43.1 (36.4 to 49.9)	<0.001
LDL <3.0 mmol/L¶	695	92.8 (90.7 to 94.5)	193	89.4 (84.5 to 93.1)	0.10
Non-HDL <2.5 mmol/L	400	52.7 (49.1 to 56.3)	89	40.6 (34.1 to 47.4)	0.002
Glucose and BP control in diabetics**	209		58		
Glucose <7 mmol/L††	61	30.4 (24.1 to 37.2)	16	29.1 (17.6 to 42.9)	0.86
HbA1c <7% (<53 mmol/mol)‡‡	107	54.9 (47.6 to 62.0)	29	50.0 (36.6 to 63.4)	0.51
HbA1c <6% (<42 mmol/mol)‡‡	29	14.9 (10.2 to 20.7)	9	15.5 (7.3 to 27.4)	0.90
BP at target‡‡	81	38.8 (32.1 to 45.7)	25	43.1 (30.2 to 56.8)	0.55

p values in bold indicate those with statistical significance

*Figures for those on blood pressure-lowering medication.

†Defined as blood pressure <140/90 mm Hg.

‡Defined as blood pressure <130/80 mm Hg.

§Figures for those on lipid-lowering medication and with cholesterol measurements made.

¶Data available for 749 male and 216 female patients.

**Figures for self-reported diabetics only.

††Data available for 201 male patients, 55 female.

‡‡Data available for 195 male patients, 58 female.

BP, blood pressure; HDL, high-density lipoprotein; LDL, high-density lipoprotein; TC, total cholesterol.

The availability and quality of these programmes should be ensured through appropriate funding and support from the Department of Health and auditing their adherence to all BACPR exemplary standards and core components. The ongoing National Audit of Cardiac Rehabilitation does not reflect all these standards and in particular there is no adequate audit of risk factors (BP, lipids and diabetes) or cardioprotective medications. NACR should consider expanding its audit function to embrace CVD as a whole and all important aspects of secondary prevention to become the National Audit for Cardiovascular Disease Prevention and Rehabilitation. The current NACR certification programme is also not comprehensive and BACPR and NACR should work together to develop performance levels which reflect the overall BACPR standards as a basis for certification.

Strengths

The main strength of A-3-P is the real-world national setting of this survey, together with the standardised methodology used to interview and examine patients using the same instruments (measuring stick, scales, tape measure and BP machine), the HemoCue point-of-care device for glucose measurement and a central laboratory for a full lipid profile, glucose, HbA1c and creatinine. All patients without self-reported diabetes were systematically investigated with an OGTT and therefore the metabolic status of all patients with CHD is described. The generic names and total daily doses of all prescribed drugs at hospital discharge and again at follow-up are recorded together with a measure of adherence.

Limitations

This survey of patients with CHD is based on a convenience sample of specialist cardiac centres and district general hospitals across the UK, with most centres located in England, followed by Scotland and fewer in Wales and Northern Ireland, which may not be representative of all hospitals. A systematic retrospective sample of consecutive patients from hospital databases was identified, but the patient response rate to interview and examination was very low, a source of potential bias. The high proportion of patients attending CR, 69% in our survey compared with 47% in the NACR database, indicates a bias towards healthier patients attending rehabilitation and therefore our results overestimate the health status of all patients with CHD and their quality of care in terms of risk factor control and drug prescribing. A truly representative patient sample is likely to include more higher risk patients with less healthier lifestyles and even poorer standards of secondary preventive care. Self-reported lifestyle has limitations but smoking habit was validated by breath carbon monoxide, and validated tools were used to measure physical activity and exercise. Diet could not be assessed using validated methods as these are too complex and time consuming and beyond the capacity of this survey.

CONCLUSION

The overall standard of preventive care remains suboptimal in the UK for patients with CHD. Although over two-thirds accessed a CR programme, inequalities still exist with less female, older adults and some diagnostic groups not attending. Despite CR attendance, lifestyle and risk factor management for several factors remains inadequate among attenders, reflecting the content and resources for these programmes. Patients require a comprehensive preventive cardiology programme with interdisciplinary expertise to address lifestyle, risk factor management and adherence to cardioprotective medications in order to achieve the standards and the specified core components set out by the BACPR and JBS3 guidelines.

Collaborators Imperial College London, National Heart and Lung Institute: D Wood*, A Adamska, S Adamska, C Jennings, K Kotseva. England Academic Cardiology, Castle Hill Hospital, Hull: A Hoye* P Atkin, D Fellowes. Barnsley Hospital NHS Foundation Trust: A Negahban*, M Cunningham, A Daniels, L Zeidan. Basildon University Hospital: Dr S Iyer*, A Nicholson, M Ocampo, A Seviliano. Blackpool Teaching Hospitals: G Galasko*, L Benham, S Preston, D Sebastian. Bradford Teaching Hospitals NHS Foundation Trust Bradford: S Lindsay*, C Atkinson, C Kranilla, M Vinod. Hammersmith Hospital: D Wood*, H Abbas, N Rhoualmi. Harrogate District NHS Foundation Trust: Y Beerachee*, C Bennett, M Broome, A Bwalya, Lindsay Caygill, L Dinning, A Gillespie, R Goodfellow, J Guy, T Idress, C Mills, C Morgan, N Oustance, N Singh, M Yare. Hinchingsbrooke Hospital: J M Jagoda*, H Bowyer, V Christenssen, A Groves. King's Mill Hospital Sherwood Forest Hospitals: A Jan Pl*, A Riaz*, M Gill, T A Sewell. Lister Hospital, Stevenage: Professor D Gorog*, M Baker, P de Sousa, T Mazenenga. London Northwest University Healthcare NHS Trust, Northwick Park Hospital, London: J Shah*, A Banfield, R Encarnado, A Taylor. Luton and Dunstable University Hospital: C Travill*, S Gent, N Hussain. North West Anglia NHS Foundation Trust: J Porter*, F Haines, T Peachey, J Taaffe, K Wells. Northumbria Healthcare NHS Foundation Trust: D P Ripley*, H Forward, H McKie, S L Pick, H E Thomas. Mid Yorkshire Hospitals NHS Trust: P D Batin*, D Exley, T Rank, J Wright. Milton Keynes University Hospital: A Kardos*, S-B Sutherland, L Wren. Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital: P Leeson*, D Barker, B Moreby, J Sawyer. Pennine Acute Hospitals NHS Trust, The Royal Oldham Hospital: J Sobolewska*, D Adams, C Corbett, K Hallett, S Kaye, L Morby, L Winstanley. Royal Berkshire Foundation NHS Trust: J Stirrup*, M Brunton. Royal Devon & Exeter Hospital, Exeter: M M Gandhi*, L Adams, A Hunt. University Hospital Coventry: L Tapp*, V Ansell, S Hyndman. University Hospitals of Morecambe Bay NHS Foundation Trust: A Brodison*, J Craig, S Peters. West Middlesex University Hospital: R Kaprielian*, A Bucaj, K Mahay, M Oblak. Wrightington Wigan and Leigh NHS Foundation Trust: A Sultan*, K Duell, M Gaskell, L Heaton, C Moore, V Parkinson, T Taylor, C Tierney, K Vandesnepscheut-Jones. Yeovil District Hospital: A Broadley*, C Buckley, L Matthews, L Pippard. York Teaching Hospital: C Gale*, M Pye*, Y McGill, H Redfearn, M Fearnley. NIHR Clinical Research Network: East Midlands, Eastern, Greater Manchester, North East and North Cumbria, North Thames, North West Coast, North West London, South West Peninsula, Thames Valley and South Midlands, West Midlands, Yorkshire and Humber. Northern Ireland: Antrim Area Hospital, Northern Ireland: K Lyons*, C Edwards, L Miskelly, S O Mullan. Royal Victoria Hospital, Belfast Health and Social Care Trust: M Spencer*, E McCart. Ulster Hospital Belfast: P Donnelly*, S Kelly, S Regan, D Turnbull. Scotland: Aberdeen Royal Infirmary: Dana Dawson*, B Brikinns, S Joseph. Golden Jubilee National Hospital: Oldroyd K*, Teyhan F, Kelly J, Tewkesbury A. Royal Infirmary of Edinburgh: D Newby*, S Clark, K Combe, L Derr, J Donnelly, L Flint, A Gill, M Glenwright, J Harrison, H Nailon, K Orr, E McDonald, C Mahoney, F Morrow, K Paterson, B Poulouse, B Rif, N Spath H Spence, A Sutherland. Raigmore Hospital: J Watt*, C Barr, S Dekker, D McDonald, L O'Keeffe, I Shread. University of Dundee: C C Lang*, A M Choy, L Douglas, S Kalra, M Mohan, R Symon. Wales: Abertawe Bro Morgannwg University Health Board: J Halcox*, K Baldwin, H Goldring, C Thomas. Prince Philip Hospital: L Izzat*, D Evans, Z Omar, E Perkins. (* indicates the principal investigator)

Contributors All authors contributed to the design, conduct, analysis and interpretation of results and take full responsibility as authors for the manuscript.

Funding This survey was funded by research grants from Amgen, AstraZeneca, Novartis, Pfizer and Sanofi through the European Society of Cardiology and the Preventive Cardiology Trust (Registered UK Charity).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethics approval was obtained from the London Central Research Ethics Committee in Spring 2016 (REC reference number 16/LO/0432).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID ID

Catriona S Jennings <http://orcid.org/0000-0002-9143-1425>

REFERENCES

- Bowker TJ, Clayton TC, Ingham J, *et al*. A British cardiac Society survey of the potential for the secondary prevention of coronary disease: ASPIRE (action on secondary prevention through intervention to reduce events). *Heart* 1996;75:334–42.
- Kotseva K, Jennings CS, Turner EL, *et al*. ASPIRE-2-PREVENT: a survey of lifestyle, risk factor management and cardioprotective medication in patients with coronary heart disease and people at high risk of developing cardiovascular disease in the UK. *Heart* 2012;98:865–71.
- Boon N, Boyle R, Bradbury K, *et al*. Joint British societies' consensus recommendations for the prevention of cardiovascular disease (JBS3). *Heart* 2014;100 Suppl 2:ii1–67.
- EUROASPIRE Study Group. EUROASPIRE: a European Society of cardiology survey of secondary prevention of coronary heart disease: principal results. *Eur Heart J* 1997;18:1569–82.
- EUROASPIRE II Study Group. Lifestyle and risk factor management and use of drug therapies in coronary patients from 15 countries; principal results from EUROASPIRE II Euro heart survey programme. *Eur Heart J* 2001;22:554–72.
- Kotseva K, Wood D, De Backer G, *et al*. EUROASPIRE III: a survey on the lifestyle, risk factors and use of cardioprotective drug therapies in coronary patients from 22 European countries. *Eur J Cardiovasc Prev Rehabil* 2009;16:121–37.
- Kotseva K, Wood D, De Bacquer D, *et al*. EUROASPIRE IV: a European Society of cardiology survey on the lifestyle, risk factor and therapeutic management of coronary patients from 24 European countries. *Eur J Prev Cardiol* 2016;23:636–48.
- Kotseva K, De Backer G, De Bacquer D, *et al*. Lifestyle and impact on cardiovascular risk factor control in coronary patients across 27 countries: results from the European Society of cardiology ESC-EORP EUROASPIRE V registry. *Eur J Prev Cardiol* 2019;26:824–35.
- Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease. *JAMA* 2003;290:86–97.
- Anderson L, Taylor RS. Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews. *Cochrane Database Syst Rev* 2014;CD011273.
- Chow CK, Jolly S, Rao-Melacini P, *et al*. Association of diet, exercise, and smoking modification with risk of early cardiovascular events after acute coronary syndromes. *Circulation* 2010;121:750–8.
- BACPR. *The BACPR standards and core components for cardiovascular disease prevention and rehabilitation 2017*. 3rd edn, 2017. http://www.bacpr.com/resources/BACPR_Standards_and_Core_Components_2017.pdf
- The National audit of cardiac rehabilitation annual statistical report, 2018. Available: <https://www.bhf.org.uk/informationsupport/publications/statistics/national-audit-of-cardiac-rehabilitation-quality-and-outcomes-report-2018>
- Whelton PK, Carey RM, Aronow WS, *et al*. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: Executive summary: a report of the American College of Cardiology/American heart association Task force on clinical practice guidelines. *Hypertension* 2018;71:1269–324.
- Williams B, Mancia G, Spiering W, *et al*. 2018 ESC/ESH guidelines for the management of arterial hypertension. *Eur Heart J* 2018;39:3021–104.
- Cahill K, Stevens S, Perera R, *et al*. Pharmacological interventions for smoking cessation: an overview and network meta-analysis. *Cochrane Database Syst Rev* 2013;5:CD009329.
- Rigotti NA, Munafò MR, Stead LF. Smoking cessation interventions for hospitalized smokers: a systematic review. *Arch Intern Med* 2008;168:1950–60.
- Jennings C, Kotseva K, De Bacquer D, *et al*. Effectiveness of a preventive cardiology programme for high CVD risk persistent smokers: the EUROACTION plus varenicline trial. *Eur Heart J* 2014;35:1411–20.
- Anthenelli RM, Benowitz NL, West R, *et al*. Neuropsychiatric safety and efficacy of varenicline, bupropion, and nicotine patch in smokers with and without psychiatric disorders (eagles): a double-blind, randomised, placebo-controlled clinical trial. *Lancet* 2016;387:2507–20.
- NHS. Long term plan, 2019. Available: <https://www.longtermplan.nhs.uk/wp-content/uploads/2019/01/nhs-long-term-plan.pdf>
- Aubin H-J, Farley A, Lycett D, *et al*. Weight gain in smokers after quitting cigarettes: meta-analysis. *BMJ* 2012;345:e4439.
- Gyberg V, De Bacquer D, Kotseva K, *et al*. Screening for dysglycaemia in patients with coronary artery disease as reflected by fasting glucose, oral glucose tolerance test, and HbA1c: a report from EUROASPIRE IV—a survey from the European Society of Cardiology. *Eur Heart J* 2015;36:1171–7.
- van Halewijn G, Deckers J, Tay HY, *et al*. Lessons from contemporary trials of cardiovascular prevention and rehabilitation: a systematic review and meta-analysis. *Int J Cardiol* 2017;232:294–303.
- Connolly SB, Kotseva K, Jennings C, *et al*. Outcomes of an integrated community-based nurse-led cardiovascular disease prevention programme. *Heart* 2017;103:840–7.
- Gibson I, Flaherty G, Cormican S, *et al*. Translating guidelines to practice: findings from a multidisciplinary preventive cardiology programme in the West of Ireland. *Eur J Prev Cardiol* 2014;21:366–76.
- WHO. *Obesity: preventing and managing the global epidemic. Report of a WHO consultation*. Geneva: World Health Organisation, 2004. [http://whqlibdoc.who.int/trs/WHO_TRS_894_\(part1\).pdf](http://whqlibdoc.who.int/trs/WHO_TRS_894_(part1).pdf)
- Inoue S, Zimmet P. The Asia-Pacific perspective. redefining obesity and its treatment, 2000. Available: http://www.diabetes.com/pdf/obesity_report.pdf
- Godin G, Shephard RJ. Godin leisure-time exercise questionnaire. *Med Sci Sports Exerc* 1997;29:S36e8.
- NICE. Managing diabetes in patients with type 2 diabetes, 2019. Available: <http://pathways.nice.org.uk/pathways/type-2-diabetes-in-adults>
- WHO. Definition and diagnosis of diabetes mellitus and intermediate hyperglycaemia [Internet], 2019. Available: https://www.who.int/diabetes/publications/diagnosis_diabetes2006/en/ [Accessed 12 Jun 2019].
- Gehi A, Haas D, Pipkin S, *et al*. Depression and medication adherence in outpatients with coronary heart disease: findings from the heart and soul study. *Arch Intern Med* 2005;165:2508–13.
- Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
- Oldridge N, Höfer S, McGee H, *et al*. The HeartQoL: Part II. validation of a new core health-related quality of life questionnaire for patients with ischemic heart disease. *Eur J Prev Cardiol* 2014;21:98–106.
- Schweikert B, Hahmann H, Leidl R. Validation of the EuroQoL questionnaire in cardiac rehabilitation. *Heart* 2006;92:62–7.