	Sacubitril/valsartan		Control			Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Random, 95% Cl	M-H, Random, 95% CI
5.4.1 Chronic							
AWAKE-HF 2019	8	69	26	70	6.4%	0.31 [0.15, 0.64]	
Fan JF 2019	2	25	2	27	1.1%	1.08 [0.16, 7.10]	-
Gao 2019	2	17	4	17	1.6%	0.50 [0.11, 2.38]	· ·
Ke ZF 2019	9	35	12	35	6.3%	0.75 [0.36, 1.55]	<del></del>
Li J 2019 (1)	1	62	1	64	0.5%	1.03 [0.07, 16.14]	
Liu YH 2019	3	26	8	22	2.6%	0.32 [0.10, 1.05]	<del></del>
OUTSTEP-HF	168	309	143	310	28.6%	1.18 [1.01, 1.38]	<del>-</del>
PARADIGM-HF 2014	2479	4203	2594	4229	34.4%	0.96 [0.93, 1.00]	
Subtotal (95% CI)		4746		4774	81.3%	0.86 [0.67, 1.10]	•
Total events	2672		2790				
Heterogeneity: Tau <sup>2</sup> = 0.0	04; Chi <sup>2</sup> = 20.1	6, df = 7	P = 0.00	5); l² =	65%		
Test for overall effect: Z =	= 1.18 (P = 0.2	4)					
5.4.2 Acute							
PIONEER-HF 2018	66	440	65	441	18.7%	1.02 [0.74, 1.40]	<del>-</del>
Subtotal (95% CI)		440		441	18.7%	1.02 [0.74, 1.40]	<b>*</b>
Total events	66		65				
Heterogeneity: Not applic	able						
Test for overall effect: Z =	= 0.11 (P = 0.9	1)					
Total (95% CI)		5186		5215	100.0%	0.91 [0.74, 1.11]	•
Total events	2738		2855				
Heterogeneity: Tau <sup>2</sup> = 0.0		5. df = 8 i	P = 0.009	9);  ² =	60%		<u> </u>
Test for overall effect: Z = 0.93 (P = 0.35)							0.05 0.2 1 5 20
Test for subgroup differences: $Chi^2 = 0.66$ , $df = 1$ (P = 0.42), $I^2 = 0\%$							Sacubitril/valsartan Control