

Supplemental Material

Supplemental Table 1. Detailed locations of PCI lesions

Supplemental Table 2. Patient and treatment status at 9 months

Supplemental Table 3. Adjusted odds ratios for the primary endpoint

Supplemental Table 1. Detailed locations of PCI lesions

Lesion		BB (N = 26)	CCB (N = 26)
RCA	#1		1
	#2		0
	#3	7	2
	#4AV		0
	#4PD		0
LAD	#5		0
	#6		11
	#7	18	6
	#8		21
	#9		2
	#10		0
LCX	#11		1
	#12		1
	#13	5	8
	#14		1
	#15		0

RCA= right coronary artery; LAD=left anterior descending; LCX=left circumflex;
 N=number of patients; AV=atrioventricular; PD= posterior descending

Supplemental Table 2. Patient and treatment status at 9 months

Number of patients	Overall		BB		CCB		<i>P</i> value
	52		26		26		
Systolic blood pressure (mmHg) mean (SD)	125	18.4	122	18.4	128	18.3	0.28
Diastolic blood pressure (mmHg) mean (SD)	75	12.5	72	12.9	78	11.5	0.07
Heart rate (bpm)	72	12.3	67	12.0	76	11.0	0.007
Beta-blocker	52%	27	93%	24	12%	3	
Carvedilol	1.9%	1	3.9%	1	3.9%	1	<0.001
Bisoprolol	50%	26	89%	23	7.7%	2	
Calcium-channel blocker*	54%	28	15%	4	92%	24	<0.001

Values are mean _ SD or n (%).

SD=standard deviation; bpm=beat per minute

* = Amlodipine

Supplemental Table 3. Adjusted odds ratios for the primary endpoint

	CCB (N = 26)	BB (N = 26)
		Odds ratio (95% CI)
Model 1	Reference	0.670 [-4.30, 3.49]
Previous MI	Reference	3.20 * 10 ^{^(-8)} [-inf, inf]
Previous PCI	Reference	2.00 [-8.09, 9.48]
Previous CABG	Reference	0.735 [-inf, inf]
Model 2	Reference	0.632 [-0.435, 3.43]
Previous PCI	Reference	0.500[-7.36, 5.98]

Inf= infinite; CCB=calcium channel blocker; BB= beta blocker; N=number of patients