

Online Table S2. Univariable analysis for mortality.

Risk factor	Dead (N=293)	Alive (N=959)	HR (95% CI)	P-value
Coronary artery disease	64 (21.8%)	60 (6.3%)	3.01 (2.27 to 3.99)	<0.001
Age, years	78.3±9.4	60.6±14.7	1.10 (1.09 to 1.11)	<0.001
Age, per 5-year increase	-	-	1.62 (1.53 to 1.72)	<0.001
Female sex	86 (29.4%)	368 (38.4%)	0.67 (0.52 to 0.87)	0.002
Smoking	32 (10.9%)	65 (6.8%)	1.61 (1.14 to 2.32)	0.011
Obesity	13 (4.4%)	33 (3.4%)	1.30 (0.75 to 2.28)	0.342
Hypertension	148 (50.5%)	350 (36.5%)	1.63 (1.30 to 2.06)	<0.001
Hyperlipidemia	32 (10.9%)	61 (6.4%)	1.64 (1.14 to 2.38)	0.008
Diabetes mellitus	57 (19.5%)	134 (14.0%)	1.38 (1.03 to 1.86)	0.030
Chronic kidney disease	41 (14.0%)	63 (6.6%)	2.00 (1.43 to 2.81)	<0.001
Prior CVE	33 (11.3%)	70 (7.3%)	1.44 (0.99 to 2.11)	0.057
LVEF <35%	17 (5.8%)	21 (2.2%)	2.13 (1.29 to 3.53)	0.003
Atrial fibrillation	37 (12.6%)	71 (7.4%)	1.68 (1.18 to 2.37)	0.004
COPD	22 (7.5%)	53 (5.5%)	1.29 (0.82 to 2.01)	0.266
Asthma	6 (2.0%)	13 (1.4%)	1.40 (0.62 to 3.13)	0.418
Active cancer	9 (3.1%)	26 (2.7%)	1.24 (0.64 to 2.40)	0.529

COPD: chronic obstructive pulmonary disease; CVE: cerebrovascular event; LVEF: left ventricular ejection fraction.