

Supplemental Figure Legends

Supplemental Figure 1. PARIS and CREDO-Kyoto thrombotic and bleeding risk scoring systems

BMI: body mass index, CrCl: creatinine clearance, CKD: chronic kidney disease, PCI: percutaneous coronary intervention, CABG: coronary bypass grafting, ACS: acute coronary syndrome, Hb: hemoglobin, CTO: chronic total occlusion.


Supplemental Figure 2. Distribution of the risk scores according to HASBRED and CHADS2-VASc risk scoring systems

Supplemental Figure 3. Distribution of the risk score categories according to PARIS and CREDO-Kyoto risk scoring systems

Supplemental Figure 1.

	Risk scores for bleeding events	Risk scores for ischemic events
PARIS score	Age, years <50: 0, 50-59: +1, 60-69: +2, 70-79: +3, ≥80: +4	Diabetes mellitus None: 0, Non-insulin: +1, insulin-dependent: +3
Ischaemic risks	BMI, kg/m ² <25: +2, 25-34.9: 0, ≥35: +2	Current smoking Yes: +1, No: 0
Low: 0-2	Current smoking Yes: +2, No: 0	CrCl < 60ml/min Present: +2, Absent: 0
Intermediate: 3-4	Anaemia Present: +3, Absent: 0	Prior PCI Yes: +2, No: 0
High: ≥5	CrCl <60 ml/min Present: +2, Absent: 0	Prior CABG Yes: +2, No: 0
Bleeding risks	Triple therapy on discharge Yes: +2, No: 0	ACS presentation is not included
Low: 0-3	Low platelet (<100000/uL): +2	Severe CKD: +2
Intermediate: 4-7	Severe CKD: +2	Atrial fibrillation: +2
High risk: ≥8	Peripheral vessel disease: +2	Peripheral vessel disease: +2
CREDO-Kyoto score	Heart failure: +2	Anaemia (Hb <11 g/dl): +2
Ischaemic risks	Prior myocardial infarction: +1	Age ≥ 75 years: +1
Low: 0	Malignancy: +1	Heart failure: +1
Intermediate: 1-2	Atrial fibrillation: +1	Diabetes mellitus: +1
High: ≥3		
Bleeding risks		
Low: 0		
Intermediate: 1-2		
High risk: ≥3		CTO is not included

Supplemental Figure 2.



Supplemental Figure 3.

